

Package: EQUIVNONINF (via r-universe)

September 16, 2024

Type Package

Title Testing for Equivalence and Noninferiority

Version 1.0.2

Date 2021-06-29

Author Stefan Wellek, Peter Ziegler

Maintainer Stefan Wellek <stefan.wellek@zi-mannheim.de>

Description Making available in R the complete set of programs accompanying S. Wellek's (2010) monograph "Testing Statistical Hypotheses of Equivalence and Noninferiority. Second Edition" (Chapman&Hall/CRC).

License CC0

Depends R (>= 3.0.0), stats, BiasedUrn

NeedsCompilation no

Repository CRAN

Date/Publication 2021-07-12 13:10:02 UTC

Contents

EQUIVNONINF-package	2
bi1st	3
bi2aeq1	4
bi2aeq2	5
bi2aeq3	7
bi2by_ni_del	8
bi2by_ni_OR	10
bi2diffac	11
bi2dipow	12
bi2rlv1	14
bi2rlv2	15
bi2st	16
bi2ste1	17
bi2ste2	18

bi2ste3	19
bi2wld_ni_del	21
cf_reh_exact	22
cf_reh_midp	23
exp1st	25
fstretch	26
gofhwex	28
gofhwex_1s	29
gofind_t	30
gofsimpt	31
mawi	33
mcnasc_ni	34
mcnby_ni	35
mcnby_ni_pp	37
mcnemasc	38
mcnempow	39
mwtie_fr	40
mwtie_xy	42
postmys	43
powsign	45
pow_abe	46
po_pbibe	47
sgnrk	48
srktie_d	50
srktie_m	51
tt1st	53
tt2st	54
Index	56

EQUIVNONINF-package *Testing for equivalence and noninferiority*

Description

The package makes available in R the complete set of programs accompanying S. Wellek's (2010) monograph "Testing Statistical Hypotheses of Equivalence and Noninferiority. Second Edition" (Chapman&Hall/CRC).

Note

In order to keep execution time of all examples below the limit set by the CRAN administration, in a number of cases the function calls shown in the documentation contain specifications which are insufficient for real applications. This holds in particular true for the width `sw` of search grids, which should be chosen to be `.001` or smaller. Similarly, the maximum number of interval halving steps to be carried out in finding maximally admissible significance levels should be set to values ≥ 10 .

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

Maintainer: Stefan Wellek <stefan.wellek@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2015.

Examples

```
bi2ste1(397, 397, 0.0, 0.025, 0.511, 0.384)
bi2ste2(0.0, 0.025, 0.95, 0.8, 0.80, 1.0)
```

bi1st	<i>Critical constants and power of the UMP test for equivalence of a single binomial proportion to some given reference value</i>
-------	---

Description

The function computes the critical constants defining the uniformly most powerful (randomized) test for the problem $p \leq p_1$ or $p \geq p_2$ versus $p_1 < p < p_2$, with p denoting the parameter of a binomial distribution from which a single sample of size n is available. In the output, one also finds the power against the alternative that the true value of p falls on the midpoint of the hypothetical equivalence interval (p_1, p_2) .

Usage

```
bi1st(alpha, n, P1, P2)
```

Arguments

alpha	significance level
n	sample size
P1	lower limit of the hypothetical equivalence range for the binomial parameter p
P2	upper limit of the hypothetical equivalence range for p

Value

alpha	significance level
n	sample size
P1	lower limit of the hypothetical equivalence range for the binomial parameter p
P2	upper limit of the hypothetical equivalence range for p

C1	left-hand limit of the critical interval for the observed number X of successes
C2	right-hand limit of the critical interval for X
GAM1	probability of rejecting the null hypothesis when it turns out that $X = C_1$
GAM2	probability of rejecting the null hypothesis for $X = C_2$
POWNONRD	Power of the nonrandomized version of the test against the alternative $p = (p_1 + p_2)/2$
POW	Power of the randomized UMP test against the alternative $p = (p_1 + p_2)/2$

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 4.3.

Examples

```
bi1st(.05,273,.65,.75)
```

 bi2aeq1

Power of the exact Fisher type test for equivalence

Description

The function computes exact values of the power of the randomized UMPU test for equivalence in the strict (i.e. two-sided) sense of two binomial distributions and the conservative nonrandomized version of that test. It is assumed that the samples being available from both distributions are independent.

Usage

```
bi2aeq1(m,n,rho1,rho2,alpha,p1,p2)
```

Arguments

m	size of Sample 1
n	size of Sample 2
rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
p1	true success rate in Population 1
p2	true success rate in Population 2

Value

m	size of Sample 1
n	size of Sample 2
rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
p1	true success rate in Population 1
p2	true success rate in Population 2
POWNR	Power of the nonrandomized version of the test
POW	Power of the randomized UMPU test

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.4.

Examples

```
bi2aeq1(302,302,0.6667,1.5,0.05,0.5,0.5)
```

 bi2aeq2

Sample sizes for the exact Fisher type test for equivalence

Description

The function computes minimum sample sizes required in the randomized UMPU test for equivalence of two binomial distributions with respect to the odds ratio. Computation is done under the side condition that the ratio m/n has some predefined value λ .

Usage

```
bi2aeq2(rho1, rho2, alpha, p1, p2, beta, qlambda)
```

Arguments

rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
p1	true success rate in Population 1
p2	true success rate in Population 2
beta	target value of power
qlambda	sample size ratio m/n

Value

rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
p1	true success rate in Population 1
p2	true success rate in Population 2
beta	target value of power
qlambda	sample size ratio m/n
M	minimum size of Sample 1
N	minimum size of Sample 2
POW	Power of the randomized UMPU test attained with the computed values of m,n

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.4.

Examples

```
bi2aeq2(0.5, 2.0, 0.05, 0.5, 0.5, 0.60, 1.0)
```

bi2aeq3	<i>Determination of a maximally raised nominal significance level for the nonrandomized version of the exact Fisher type test for equivalence</i>
---------	---

Description

The objective is to raise the nominal significance level as far as possible without exceeding the target significance level in the nonrandomized version of the test. The approach goes back to R.D. Boschloo (1970) who used the same technique for reducing the conservatism of the traditional nonrandomized Fisher test for superiority.

Usage

```
bi2aeq3(m,n,rho1,rho2,alpha,sw,tolrd,tol,maxh)
```

Arguments

m	size of Sample 1
n	size of Sample 2
rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
tolrd	horizontal distance from 0 and 1, respectively of the left- and right-most boundary point to be included in the search grid
tol	upper bound to the absolute difference between size and target level below which the search for a corrected nominal level terminates
maxh	maximum number of interval halving steps to be carried out in finding the maximally raised nominal level

Details

It should be noted that, as the function of the nominal level, the size of the nonrandomized test is piecewise constant. Accordingly, there is a nondegenerate interval of "candidate" nominal levels serving the purpose. The limits of such an interval can be read from the output. In terms of execution time, bi2aeq3 is the most demanding program of the whole package.

Value

m	size of Sample 1
n	size of Sample 2
rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio

alpha	significance level
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
tolrd	horizontal distance from 0 and 1, respectively of the left- and right-most boundary point to be included in the search grid
tol	upper bound to the absolute difference between size and target level below which the search for a corrected nominal level terminates
maxh	maximum number of interval halving steps to be carried out in finding the maximally raised nominal level
ALPH_0	current trial value of the raised nominal level searched for
NHST	number of interval-halving steps performed up to now
SIZE	size of the critical region corresponding to α_0

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Boschloo RD: Raised conditional level of significance for the 2 x 2- table when testing the equality of two probabilities. *Statistica Neerlandica* 24 (1970), 1-35.

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.5.

Examples

```
bi2aeq3(50,50,0.6667,1.5000,0.05,0.01,0.000001,0.0001,5)
```

bi2by_ni_del	<i>Objective Bayesian test for noninferiority in the two-sample setting with binary data and the difference of the two proportions as the parameter of interest</i>
--------------	---

Description

Implementation of the construction described on pp. 185-6 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

```
bi2by_ni_del(N1, N2, EPS, SW, NSUB, ALPHA, MAXH)
```


Arguments

N1	size of Sample 1
N2	size of sample 2
EPS	noninferiority margin to the difference of success probabilities
SW	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
NSUB	number of subintervals for partitioning the range of integration
ALPHA	target significance level
MAXH	maximum number of interval halving steps to be carried out in finding the maximally admissible nominal level

Details

The program uses 96-point Gauss-Legendre quadrature on each of the NSUB intervals into which the range of integration is partitioned.

Value

N1	size of Sample 1
N2	size of sample 2
EPS	noninferiority margin to the difference of success probabilities
NSUB	number of subintervals for partitioning the range of integration
SW	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
ALPHA0	result of the search for the largest admissible nominal level
SIZE0	size of the critical region corresponding to α_0
SIZE_UNC	size of the critical region of the test at uncorrected nominal level

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Statistical methods for the analysis of two-armed non-inferiority trials with binary outcomes. *Biometrical Journal* 47 (2005), 48–61.

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.3.

Examples

```
bi2by_ni_del(20,20,.10,.01,10,.05,5)
```

 bi2by_ni_OR

Objective Bayesian test for noninferiority in the two-sample setting with binary data and the odds ratio as the parameter of interest

Description

Implementation of the construction described on pp. 179–181 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

bi2by_ni_OR(N1, N2, EPS, SW, NSUB, ALPHA, MAXH)

Arguments

N1	size of sample 1
N2	size of sample 2
EPS	noninferiority margin to the deviation of the odds ratio from unity
SW	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
NSUB	number of subintervals for partitioning the range of integration
ALPHA	target significance level
MAXH	maximum number of interval halving steps to be carried out in finding the maximally admissible nominal level

Details

The program uses 96-point Gauss-Legendre quadrature on each of the NSUB intervals into which the range of integration is partitioned.

Value

N1	size of sample 1
N2	size of sample 2
EPS	noninferiority margin to the deviation of the odds ratio from unity
NSUB	number of subintervals for partitioning the range of integration
SW	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
ALPHA0	result of the search for the largest admissible nominal level
SIZE0	size of the critical region corresponding to α_0
SIZE_UNC	size of the critical region of the test at uncorrected nominal level

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Statistical methods for the analysis of two-arm non-inferiority trials with binary outcomes. *Biometrical Journal* 47 (2005), 48–61.
 Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.2.

Examples

```
bi2by_ni_OR(10,10,1/3,.0005,10,.05,12)
```

bi2diffac	<i>Determination of a corrected nominal significance level for the asymptotic test for equivalence of two unrelated binomial proportions with respect to the difference δ of their population counterparts</i>
-----------	--

Description

The program computes the largest nominal significance level which can be substituted for the target level α without making the exact size of the asymptotic testing procedure larger than α .

Usage

```
bi2diffac(alpha,m,n,del1,del2,sw,tolrd,tol,maxh)
```

Arguments

alpha	significance level
m	size of Sample 1
n	size of Sample 2
del1	absolute value of the lower limit of the hypothetical equivalence range for $p_1 - p_2$
del2	upper limit of the hypothetical equivalence range for $p_1 - p_2$
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
tolrd	horizontal distance of the left- and right-most boundary point to be included in the search grid
tol	upper bound to the absolute difference between size and target level below which the search for a corrected nominal level terminates
maxh	maximum number of interval halving steps to be carried out in finding the maximally raised nominal level

Value

alpha	significance level
m	size of Sample 1
n	size of Sample 2
de11	absolute value of the lower limit of the hypothetical equivalence range for $p_1 - p_2$
de12	upper limit of the hypothetical equivalence range for $p_1 - p_2$
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
tolrd	horizontal distance of the left- and right-most boundary point to be included in the search grid
tol	upper bound to the absolute difference between size and target level below which the search for a corrected nominal level terminates
maxh	maximum number of interval halving steps to be carried out in finding the maximally raised nominal level
NH	number of interval-halving steps actually performed
ALPH_0	value of the raised nominal level obtained after NH steps
SIZE0	size of the critical region corresponding to α_0
ERROR	error indicator answering the question of whether or not the sufficient condition for the correctness of the result output by the program, was satisfied

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.6.

Examples

```
bi2diffac(0.05,20,20,0.40,0.40,0.1,1e-6,1e-4,3)
```

 bi2dipow

Exact rejection probability of the asymptotic test for equivalence of two unrelated binomial proportions with respect to the difference of their expectations at any nominal level under an arbitrary parameter configuration

Description

The program computes exact values of the rejection probability of the asymptotic test for equivalence in the sense of $-\delta_1 < p_1 - p_2 < \delta_2$, at any nominal level α_0 . [The largest α_0 for which the test is valid in terms of the significance level, can be computed by means of the program bi2diffac.]

Usage

```
bi2dipow(alpha0,m,n,de11,de12,p1,p2)
```

Arguments

alpha0	nominal significance level
m	size of Sample 1
n	size of Sample 2
de11	absolute value of the lower limit of the hypothetical equivalence range for $p_1 - p_2$
de12	upper limit of the hypothetical equivalence range for $p_1 - p_2$
p1	true value of the success probability in Population 1
p2	true value of the success probability in Population 2

Value

alpha0	nominal significance level
m	size of Sample 1
n	size of Sample 2
de11	absolute value of the lower limit of the hypothetical equivalence range for $p_1 - p_2$
de12	upper limit of the hypothetical equivalence range for $p_1 - p_2$
p1	true value of the success probability in Population 1
p2	true value of the success probability in Population 2
POWEX0	exact rejection probability under (p_1, p_2) of the test at nominal level α_0 for equivalence of two binomial distributions with respect to the difference of the success probabilities
ERROR	error indicator answering the question of whether or not the sufficient condition for the correctness of the result output by the program, was satisfied

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>

Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.6.

Examples

```
bi2dipow(0.0228,50,50,0.20,0.20,0.50,0.50)
```

 bi2rlv1

Power of the exact Fisher type test for relevant differences

Description

The function computes exact values of the power of the randomized UMPU test for relevant differences between two binomial distributions and the conservative nonrandomized version of that test. It is assumed that the samples being available from both distributions are independent.

Usage

```
bi2rlv1(m,n,rho1,rho2,alpha,p1,p2)
```

Arguments

m	size of Sample 1
n	size of Sample 2
rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
p1	true success rate in Population 1
p2	true success rate in Population 2

Value

m	size of Sample 1
n	size of Sample 2
rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
p1	true success rate in Population 1
p2	true success rate in Population 2
POWNR	power of the nonrandomized version of the test
POW	power of the randomized UMPU test

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 11.3.3.

Examples

```
bi2rlv1(200,300,.6667,1.5,.05,.25,.10)
```

bi2rlv2

Sample sizes for the exact Fisher type test for relevant differences

Description

The function computes minimum sample sizes required in the randomized UMPU test for relevant differences between two binomial distributions with respect to the odds ratio. Computation is done under the side condition that the ratio m/n has some predefined value λ .

Usage

```
bi2rlv2(rho1, rho2, alpha, p1, p2, beta, qlambda)
```

Arguments

rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
p1	true success rate in Population 1
p2	true success rate in Population 2
beta	target value of power
qlambda	sample size ratio m/n

Value

rho1	lower limit of the hypothetical equivalence range for the odds ratio
rho2	upper limit of the hypothetical equivalence range for the odds ratio
alpha	significance level
p1	true success rate in Population 1
p2	true success rate in Population 2
beta	target value of power
qlambda	sample size ratio m/n
M	minimum size of Sample 1
N	minimum size of Sample 2
POW	power of the randomized UMPU test attained with the computed values of m, n

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 11.3.3.

Examples

```
bi2rlv2(.6667,1.5,.05,.70,.50,.50,2.0)
```

bi2st	<i>Critical constants for the exact Fisher type UMPU test for equivalence of two binomial distributions with respect to the odds ratio</i>
-------	--

Description

The function computes the critical constants defining the uniformly most powerful unbiased test for equivalence of two binomial distributions with parameters p_1 and p_2 in terms of the odds ratio. Like the ordinary Fisher type test of the null hypothesis $p_1 = p_2$, the test is conditional on the total number S of successes in the pooled sample.

Usage

```
bi2st(alpha,m,n,s,rho1,rho2)
```

Arguments

alpha	significance level
m	size of Sample 1
n	size of Sample 2
s	observed total count of successes
rho1	lower limit of the hypothetical equivalence range for the odds ratio $\varrho = \frac{p_1(1-p_2)}{p_2(1-p_1)}$
rho2	upper limit of the hypothetical equivalence range for ϱ

Value

alpha	significance level
m	size of Sample 1
n	size of Sample 2
s	observed total count of successes
rho1	lower limit of the hypothetical equivalence range for the odds ratio $\varrho = \frac{p_1(1-p_2)}{p_2(1-p_1)}$
rho2	upper limit of the hypothetical equivalence range for ϱ
C1	left-hand limit of the critical interval for the number X of successes observed in Sample 1
C2	right-hand limit of the critical interval for X
GAM1	probability of rejecting the null hypothesis when it turns out that $X = C_1$
GAM2	probability of rejecting the null hypothesis for $X = C_2$

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.4.

Examples

```
bi2st(.05,225,119,171, 2/3, 3/2)
```

 bi2ste1

Power of the exact Fisher type test for noninferiority

Description

The function computes exact values of the power of the randomized UMPU test for one-sided equivalence of two binomial distributions and its conservative nonrandomized version. It is assumed that the samples being available from both distributions are independent.

Usage

```
bi2ste1(m, n, eps, alpha, p1, p2)
```

Arguments

m	size of Sample 1
n	size of Sample 2
eps	noninferiority margin to the odds ratio ϱ , defined to be the maximum acceptable deviation of the true value of ϱ from unity
alpha	significance level
p1	success rate in Population 1
p2	success rate in Population 2

Value

m	size of Sample 1
n	size of Sample 2
eps	noninferiority margin to the odds ratio ϱ , defined to be the maximum acceptable deviation of the true value of ϱ from unity
alpha	significance level
p1	success rate in Population 1
p2	success rate in Population 2
POWNR	power of the nonrandomized version of the test
POW	power of the randomized UMPU test

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, 6.6.1.

Examples

```
bi2ste1(106,107,0.5,0.05,0.9245,0.9065)
```

 bi2ste2

Sample sizes for the exact Fisher type test for noninferiority

Description

Sample sizes for the exact Fisher type test for noninferiority

Usage

```
bi2ste2(eps, alpha, p1, p2, bet, qlambda)
```

Arguments

eps	noninferiority margin to the odds ratio
alpha	significance level
p1	success rate in Population 1
p2	success rate in Population 2
bet	target power value
qlambda	sample size ratio m/n

Details

The program computes the smallest sample sizes m, n satisfying $m/n = \lambda$ required for ensuring that the power of the randomized UMPU test does not fall below β .

Value

eps	noninferiority margin to the odds ratio
alpha	significance level
p1	success rate in Population 1
p2	success rate in Population 2
bet	target power value

qlambda	sample size ratio m/n
M	minimum size of Sample 1
N	minimum size of Sample 2
POW	power of the randomized UMPU test attained with the computed values of m, n

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, 6.6.1.

Examples

```
bi2ste2(0.5, 0.05, 0.9245, 0.9065, 0.80, 1.00)
```

bi2ste3	<i>Determination of a maximally raised nominal significance level for the nonrandomized version of the exact Fisher type test for noninferiority</i>
---------	--

Description

The objective is to raise the nominal significance level as far as possible without exceeding the target significance level in the nonrandomized version of the test. The approach goes back to R.D. Boschloo (1970) who used the same technique for reducing the conservatism of the traditional nonrandomized Fisher test for superiority.

Usage

```
bi2ste3(m, n, eps, alpha, sw, tolrd, tol, maxh)
```

Arguments

m	size of Sample 1
n	size of Sample 2
eps	noninferiority margin to the odds ratio ϱ , defined to be the maximum acceptable deviation of the true value of ϱ from unity
alpha	target significance level
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
tolrd	horizontal distance from 0 and 1, respectively, of the left- and right-most boundary point to be included in the search grid

tol	upper bound to the absolute difference between size and target level below which the search for a corrected nominal level terminates
maxh	maximum number of interval-halving steps to be carried out in finding the maximally raised nominal level

Details

It should be noted that, as the function of the nominal level, the size of the nonrandomized test is piecewise constant. Accordingly, there is a nondegenerate interval of "candidate" nominal levels serving the purpose. The limits of such an interval can be read from the output.

Value

m	size of Sample 1
n	size of Sample 2
eps	noninferiority margin to the odds ratio ϱ , defined to be the maximum acceptable deviation of the true value of ϱ from unity
alpha	target significance level
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
tolrd	horizontal distance from 0 and 1, respectively, of the left- and right-most boundary point to be included in the search grid
tol	upper bound to the absolute difference between size and target level below which the search for a corrected nominal level terminates
maxh	maximum number of interval-halving steps to be carried out in finding the maximally raised nominal level
ALPH_0	current trial value of the raised nominal level searched for
NHST	number of interval-halving steps performed up to now
SIZE	size of the critical region corresponding to α_0

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

- Boschloo RD: Raised conditional level of significance for the 2 x 2- table when testing the equality of two probabilities. *Statistica Neerlandica* 24 (1970), 1-35.
- Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, §6.6.2.

Examples

```
bi2ste3(50, 50, 1/3, 0.05, 0.05, 1e-10, 1e-8, 10)
```

bi2wld_ni_del	<i>Function to compute corrected nominal levels for the Wald type (asymptotic) test for one-sided equivalence of two binomial distributions with respect to the difference of success rates</i>
---------------	---

Description

Implementation of the construction described on pp. 183-5 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

```
bi2wld_ni_del(N1, N2, EPS, SW, ALPHA, MAXH)
```

Arguments

N1	size of Sample 1
N2	size of Sample 2
EPS	noninferiority margin to the difference of success probabilities
SW	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
ALPHA	target significance level
MAXH	maximum number of interval-halving steps

Details

The program computes the largest nominal significance level to be used for determining the critical lower bound to the Wald-type statistic for the problem of testing $H : p_1 \leq p_2 - \varepsilon$ versus $K : p_1 < p_2 - \varepsilon$.

Value

N1	size of Sample 1
N2	size of Sample 2
EPS	noninferiority margin to the difference of success probabilities
SW	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
ALPHA	target significance level
MAXH	maximum number of interval-halving steps
ALPHA0	corrected nominal level
SIZE0	size of the critical region of the test at nominal level ALPHA0
SIZE_UNC	size of the test at uncorrected nominal level ALPHA
ERR_IND	indicator taking value 1 when it occurs that the sufficient condition allowing one to restrict the search for the maximum of the rejection probability under the null hypothesis to its boundary, fails to be satisfied; otherwise the indicator retains its default value 0.

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.6.3.

Examples

```
bi2wld_ni_del(25,25,.10,.01,.05,10)
```

cf_reh_exact	<i>Exact confidence bounds to the relative excess heterozygosity (REH) exhibited by a SNP genotype distribution</i>
--------------	---

Description

Implementation of the interval estimation procedure described on pp. 305-6 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

```
cf_reh_exact(X1,X2,X3,alpha,SW,TOL,ITMAX)
```

Arguments

X1	count of homozygotes of the first kind [\leftrightarrow genotype AA]
X2	count of heterozygotes [\leftrightarrow genotype AB]
X3	count of homozygotes of the second kind [\leftrightarrow genotype BB]
alpha	1 - confidence level
SW	width of the search grid for determining an interval covering the parameter point at which the conditional distribution function takes value α and $1 - \alpha$, respectively
TOL	numerical tolerance to the deviation between the computed confidence limits and their exact values
ITMAX	maximum number of interval-halving steps

Details

The program exploits the structure of the family of all genotype distributions, which is 2-parameter exponential with $\log(REH)$ as one of these parameters.

Value

X1	count of homozygotes of the first kind [\leftrightarrow genotype AA]
X2	count of heterozygotes [\leftrightarrow genotype AB]
X3	count of homozygotes of the second kind [\leftrightarrow genotype BB]
alpha	1 - confidence level
SW	width of the search grid for determining an interval covering the parameter point at which the conditional distribution function takes value α and $1 - \alpha$, respectively
TOL	numerical tolerance to the deviation between the computed confidence limits and their exact values
ITMAX	maximum number of interval-halving steps
C_l_exact	exact conditional lower $(1 - \alpha)$ -confidence bound to REH
C_r_exact	exact conditional upper $(1 - \alpha)$ -confidence bound to REH

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S, Goddard KAB, Ziegler A: A confidence-limit-based approach to the assessment of Hardy-Weinberg equilibrium. *Biometrical Journal* 52 (2010), 253-270.

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 9.4.3.

Examples

```
cf_reh_exact(34,118,96,.05,.1,1E-4,25)
```

cf_reh_midp	<i>Mid-p-value - based confidence bounds to the relative excess heterozygosity (REH) exhibited by a SNP genotype distribution</i>
-------------	---

Description

Implementation of the interval estimation procedure described on pp. 306-7 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

```
cf_reh_midp(X1,X2,X3,alpha,SW,TOL,ITMAX)
```

Arguments

X1	count of homozygotes of the first kind [\leftrightarrow genotype AA]
X2	count of heterozygotes [\leftrightarrow genotype AB]
X3	count of homozygotes of the second kind [\leftrightarrow genotype BB]
alpha	1 - confidence level
SW	width of the search grid for determining an interval covering the parameter point at which the conditional distribution function takes value α and $1 - \alpha$, respectively
TOL	numerical tolerance to the deviation between the computed confidence limits and their exact values
ITMAX	maximum number of interval-halving steps

Details

The mid-p algorithm serves as a device for reducing the conservatism inherent in exact confidence estimation procedures for parameters of discrete distributions.

Value

X1	count of homozygotes of the first kind [\leftrightarrow genotype AA]
X2	count of heterozygotes [\leftrightarrow genotype AB]
X3	count of homozygotes of the second kind [\leftrightarrow genotype BB]
alpha	1 - confidence level
SW	width of the search grid for determining an interval covering the parameter point at which the conditional distribution function takes value α and $1 - \alpha$, respectively
TOL	numerical tolerance to the deviation between the computed confidence limits and their exact values
ITMAX	maximum number of interval-halving steps
C_l_midp	lower $(1 - \alpha)$ -confidence bound to REH based on conditional mid-p-values
C_r_midp	upper $(1 - \alpha)$ -confidence bound to REH based on conditional mid-p-values

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

- Agresti A: Categorical data Analysis (2nd edn). Hoboken, NJ: Wiley, Inc., 2002, Section 1.4.5.
- Wellek S, Goddard KAB, Ziegler A: A confidence-limit-based approach to the assessment of Hardy-Weinberg equilibrium. *Biometrical Journal* 52 (2010), 253-270.
- Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 9.4.3.

Examples

```
cf_reh_midp(137,34,8,.05,.1,1E-4,25)
```

exp1st	<i>Critical constants and power against the null alternative of the UMP test for equivalence of the hazard rate of a single exponential distribution to some given reference value</i>
--------	--

Description

The function computes the critical constants defining the uniformly most powerful test for the problem $\sigma \leq 1/(1 + \varepsilon)$ or $\sigma \geq (1 + \varepsilon)$ versus $1/(1 + \varepsilon) < \sigma < (1 + \varepsilon)$, with σ denoting the scale parameter [\equiv reciprocal hazard rate] of an exponential distribution.

Usage

```
exp1st(alpha, tol, itmax, n, eps)
```

Arguments

alpha	significance level
tol	tolerable deviation from α of the rejection probability at either boundary of the hypothetical equivalence interval
itmax	maximum number of iteration steps
n	sample size
eps	margin determining the hypothetical equivalence range symmetrically on the log-scale

Value

alpha	significance level
tol	tolerable deviation from α of the rejection probability at either boundary of the hypothetical equivalence interval
itmax	maximum number of iteration steps
n	sample size
eps	margin determining the hypothetical equivalence range symmetrically on the log-scale
IT	number of iteration steps performed until reaching the stopping criterion corresponding to TOL
C1	left-hand limit of the critical interval for $T = \sum_{i=1}^n X_i$
C2	right-hand limit of the critical interval for $T = \sum_{i=1}^n X_i$
ERR1	deviation of the rejection probability from α under $\sigma = 1/(1 + \varepsilon)$
POW0	power of the randomized UMP test against the alternative $\sigma = 1$

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 4.2.

Examples

```
exp1st(0.05, 1.0e-10, 100, 80, 0.3)
```

fstretch	<i>Critical constants and power of the UMPI (uniformly most powerful invariant) test for dispersion equivalence of two Gaussian distributions</i>
----------	---

Description

The function computes the critical constants defining the optimal test for the problem $\sigma^2/\tau^2 \leq \varrho_1$ or $\sigma^2/\tau^2 \geq \varrho_2$ versus $\varrho_1 < \sigma^2/\tau^2 < \varrho_2$, with (ϱ_1, ϱ_2) as a fixed nonempty interval around unity.

Usage

```
fstretch(alpha, tol, itmax, ny1, ny2, rho1, rho2)
```

Arguments

alpha	significance level
tol	tolerable deviation from α of the rejection probability at either boundary of the hypothetical equivalence interval
itmax	maximum number of iteration steps
ny1	number of degrees of freedom of the estimator of σ^2
ny2	number of degrees of freedom of the estimator of τ^2
rho1	lower equivalence limit to σ^2/τ^2
rho2	upper equivalence limit to σ^2/τ^2

Value

alpha	significance level
tol	tolerable deviation from α of the rejection probability at either boundary of the hypothetical equivalence interval
itmax	maximum number of iteration steps
ny1	number of degrees of freedom of the estimator of σ^2

ny2	number of degrees of freedom of the estimator of τ^2
rho1	lower equivalence limit to σ^2/τ^2
rho2	upper equivalence limit to σ^2/τ^2
IT	number of iteration steps performed until reaching the stopping criterion corresponding to TOL
C1	left-hand limit of the critical interval for

$$T = \frac{n-1}{m-1} \sum_{i=1}^m (X_i - \bar{X})^2 / \sum_{j=1}^{n-1} (Y_j - \bar{Y})^2$$

C2	right-hand limit of the critical interval for
----	---

$$T = \frac{n-1}{m-1} \sum_{i=1}^m (X_i - \bar{X})^2 / \sum_{j=1}^{n-1} (Y_j - \bar{Y})^2$$

ERR	deviation of the rejection probability from α under $\sigma^2/\tau^2 = \varrho_1$
POW0	power of the UMPI test against the alternative $\sigma^2/\tau^2 = 1$

Note

If the two independent samples under analysis are from exponential rather than Gaussian distributions, the critical constants computed by means of `fstretch` with $\nu_1 = 2m$, $\nu_2 = 2n$, can be used for testing for equivalence with respect to the ratio of hazard rates. The only difference is that the ratio of sample means rather than variances has to be used as the test statistic then.

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>

Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.5.

Examples

```
fstretch(0.05, 1.0e-10, 50, 40, 45, 0.5625, 1.7689)
```

gofhwex

Critical constants of the exact UMPU test for approximate compatibility of a SNP genotype distribution with the Hardy-Weinberg model

Description

The function computes the critical constants defining the uniformly most powerful unbiased test for equivalence of the population distribution of the three genotypes distinguishable in terms of a single nucleotide polymorphism (SNP), to a distribution being in Hardy-Weinberg equilibrium (HWE).

The test is conditional on the total count S of alleles of the kind of interest, and the parameter θ , in terms of which equivalence shall be established, is defined by $\theta = \frac{\pi_2^2}{\pi_1(1-\pi_1-\pi_2)}$, with π_1 and π_2 denoting the population frequency of homozygotes of the 1st kind and heterozygotes, respectively.

Usage

```
gofhwex(alpha, n, s, de11, de12)
```

Arguments

alpha	significance level
n	number of genotyped individuals
s	observed count of alleles of the kind of interest
de11	absolute value of the lower equivalence limit to $\theta/4 - 1$
de12	upper equivalence limit to $\theta/4 - 1$

Value

alpha	significance level
n	number of genotyped individuals
s	observed count of alleles of the kind of interest
de11	absolute value of the lower equivalence limit to $\theta/4 - 1$
de12	upper equivalence limit to $\theta/4 - 1$
C1	left-hand limit of the critical interval for the observed number X_2 of heterozygotes
C2	right-hand limit of the critical interval for the observed number X_2
GAM1	probability of rejecting the null hypothesis when it turns out that $X_2 = C_1$
GAM2	probability of rejecting the null hypothesis for $X_2 = C_2$

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>

Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Tests for establishing compatibility of an observed genotype distribution with Hardy-Weinberg equilibrium in the case of a biallelic locus. *Biometrics* 60 (2004), 694-703.

Goddard KAB, Ziegler A, Wellek S: Adapting the logical basis of tests for Hardy-Weinberg equilibrium to the real needs of association studies in human and medical genetics. *Genetic Epidemiology* 33 (2009), 569-580.

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 9.4.2.

Examples

```
gofhwex(0.05, 475, 429, 1-1/1.96, 0.96)
```

gofhwex_1s	<i>Critical constants of the exact UMPU test for absence of a substantial deficit of heterozygotes as compared with a HWE-compliant SNP genotype distribution [noninferiority version of the test implemented by means of gofhwex]</i>
------------	--

Description

The function computes the critical constants defining the UMPU test for one-sided equivalence of the population distribution of a SNP, to a distribution being in Hardy-Weinberg equilibrium (HWE). A substantial deficit of heterozygotes is defined to occur when the true value of the parametric function $\omega = \frac{\pi_2/2}{\sqrt{\pi_1\pi_3}}$ [called relative excess heterozygosity (REH)] falls below unity by more than some given margin δ_0 .

Like its two-sided counterpart [see the description of the R function gofhwex], the test is conditional on the total count S of alleles of the kind of interest.

Usage

```
gofhwex_1s(alpha, n, s, del0)
```

Arguments

alpha	significance level
n	number of genotyped individuals
s	observed count of alleles of the kind of interest
del0	noninferiority margin for ω , which has to satisfy $\omega > 1 - \delta_0$ under the alternative hypothesis to be established

Value

alpha	significance level
n	number of genotyped individuals
s	observed count of alleles of the kind of interest
de10	noninferiority margin for ω , which has to satisfy $\omega > 1 - \delta_0$ under the alternative hypothesis to be established
C	left-hand limit of the critical interval for the observed number X_2 of heterozygotes
GAM	probability of rejecting the null hypothesis when it turns out that $X_2 = C$

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, pp. 300-302.

Examples

```
gofhwex_1s(0.05, 133, 65, 1-1/1.96)
```

gofind_t	<i>Establishing approximate independence in a two-way contingency table: Test statistic and critical bound</i>
----------	--

Description

The function computes all quantities required for carrying out the asymptotic test for approximate independence of two categorical variables derived in § 9.2 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

```
gofind_t(alpha, r, s, eps, xv)
```

Arguments

alpha	significance level
r	number of rows of the contingency table under analysis
s	number of columns of the contingency table under analysis
eps	margin to the Euclidean distance between the vector π of true cell probabilities and the associated vector of products of marginal totals
xv	row vector of length $r \cdot s$ whose $(i - 1)s + j$ -th component is the entry in cell (i, j) of the $r \times s$ contingency table under analysis $i = 1, \dots, r, j = 1, \dots, s$.

Value

n	size of the sample to which the input table relates
alpha	significance level
r	number of rows of the contingency table under analysis
s	number of columns of the contingency table under analysis
eps	margin to the Euclidean distance between the vector π of true cell probabilities and the associated vector of products of marginal totals
X(r, s)	observed cell counts
DSQ_OBS	observed value of the squared Euclidean distance
VN	square root of the estimated asymptotic variance of $\sqrt{n}DSQ_OBS$
CRIT	upper critical bound to $\sqrt{n}DSQ_OBS$
REJ	indicator of a positive [=1] vs negative [=0] rejection decision to be taken with the data under analysis

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>

Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 9.2.

Examples

```
xv <- c(8, 13, 15, 6, 19, 21, 31, 7)
gofind_t(0.05, 2, 4, 0.15, xv)
```

gofsimpt

Establishing goodness of fit of an observed to a fully specified multinomial distribution: test statistic and critical bound

Description

The function computes all quantities required for carrying out the asymptotic test for goodness rather than lack of fit of an observed to a fully specified multinomial distribution derived in § 9.1 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

```
gofsimpt(alpha, n, k, eps, x, pio)
```

Arguments

alpha	significance level
n	sample size
k	number of categories
eps	margin to the Euclidean distance between the vectors π and π_0 of true and hypothesized cell probabilities
x	vector of length k with the observed cell counts as components
pio	prespecified vector of cell probabilities

Value

alpha	significance level
n	sample size
k	number of categories
eps	margin to the Euclidean distance between the vectors π and π_0 of true and hypothesized cell probabilities
X(1, K)	observed cell counts
PI0(1, K)	hypothesized cell probabilities
DSQPIH_0	observed value of the squared Euclidean distance
VN_N	square root of the estimated asymptotic variance of $\sqrt{n}DSQPIH_0$
CRIT	upper critical bound to $\sqrt{n}DSQPIH_0$
REJ	indicator of a positive [=1] vs negative [=0] rejection decision to be taken with the data under analysis

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 9.1.

Examples

```
x<- c(17,16,25,9,16,17)
pio <- rep(1,6)/6
gofsimpt(0.05,100,6,0.15,x,pio)
```

mawi	<i>Mann-Whitney test for equivalence of two continuous distributions of arbitrary shape: test statistic and critical upper bound</i>
------	--

Description

Implementation of the asymptotically distribution-free test for equivalence of two continuous distributions in terms of the Mann-Whitney-Wilcoxon functional. For details see Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition, § 6.2.

Usage

```
mawi(alpha,m,n,eps1_,eps2_,x,y)
```

Arguments

alpha	significance level
m	size of Sample 1
n	size of Sample 2
eps1_	absolute value of the left-hand limit of the hypothetical equivalence range for $\pi_+ - 1/2$
eps2_	right-hand limit of the hypothetical equivalence range for $\pi_+ - 1/2$
x	row vector with the m observations making up Sample1 as components
y	row vector with the n observations making up Sample2 as components

Details

Notation: π_+ stands for the Mann-Whitney functional defined by $\pi_+ = P[X > Y]$, with $X \sim F \equiv$ cdf of Population 1 being independent of $Y \sim G \equiv$ cdf of Population 2.

Value

alpha	significance level
m	size of Sample 1
n	size of Sample 2
eps1_	absolute value of the left-hand limit of the hypothetical equivalence range for $\pi_+ - 1/2$
eps2_	right-hand limit of the hypothetical equivalence range for $\pi_+ - 1/2$
W+	observed value of the U -statistics estimator of π_+
SIGMAH	square root of the estimated asymptotic variance of W_+
CRIT	upper critical bound to $ W_+ - 1/2 - (\varepsilon'_2 - \varepsilon'_1)/2 /\hat{\sigma}$
REJ	indicator of a positive [=1] vs negative [=0] rejection decision to be taken with the data under analysis

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: A new approach to equivalence assessment in standard comparative bioavailability trials by means of the Mann-Whitney statistic. *Biometrical Journal* 38 (1996), 695-710.

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.2.

Examples

```
x <- c(10.3, 11.3, 2.0, -6.1, 6.2, 6.8, 3.7, -3.3, -3.6, -3.5, 13.7, 12.6)
y <- c(3.3, 17.7, 6.7, 11.1, -5.8, 6.9, 5.8, 3.0, 6.0, 3.5, 18.7, 9.6)
mawi(0.05, 12, 12, 0.1382, 0.2602, x, y)
```

 mcnasc_ni

Determination of a corrected nominal significance level for the asymptotic test for noninferiority in the McNemar setting

Description

The program computes the largest nominal significance level which can be substituted for the target level α without making the exact size of the asymptotic testing procedure larger than α .

Usage

```
mcnasc_ni(alpha, n, del0, sw, tol, maxh)
```

Arguments

alpha	significance level
n	sample size
del0	absolute value of the noninferiority margin for $\delta := p_{10} - p_{01}$, with p_{10} and p_{01} denoting the probabilities of discordant pairs of both kinds
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
tol	upper bound to the absolute difference between size and target level below which the search for a corrected nominal level terminates
maxh	maximum number of interval halving steps to be carried out in finding the maximally raised nominal level

Value

alpha	significance level
n	sample size
del0	absolute value of the noninferiority margin for $\delta := p_{10} - p_{01}$, with p_{10} and p_{01} denoting the probabilities of discordant pairs of both kinds
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
ALPH_0	value of the corrected nominal level obtained after nh steps
SIZE_UNC	exact size of the rejection region of the test at uncorrected nominal level α
SIZE0	exact size of the rejection region of the test at nominal level α_0
NH	number of interval-halving steps actually performed

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>

Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 5.2.3.

Examples

```
mcnasc_ni(0.05, 50, 0.05, 0.05, 0.0001, 5)
```

mcnby_ni	<i>Bayesian test for noninferiority in the McNemar setting with the difference of proportions as the parameter of interest</i>
----------	--

Description

The program determines through iteration the largest nominal level α_0 such that comparing the posterior probability of the alternative hypothesis $K_1 : \delta > -\delta_0$ to the lower bound $1 - \alpha_0$ generates a critical region whose size does not exceed the target significance level α . In addition, exact values of the power against specific parameter configurations with $\delta = 0$ are output.

Usage

```
mcnby_ni(N, DEL0, K1, K2, K3, NSUB, SW, ALPHA, MAXH)
```

Arguments

N	sample size
DEL0	noninferiority margin to the difference of the parameters of the marginal binomial distributions under comparison
K1	Parameter 1 of the Dirichlet prior for the family of trinomial distributions
K2	Parameter 2 of the Dirichlet prior for the family of trinomial distributions
K3	Parameter 3 of the Dirichlet prior for the family of trinomial distributions
NSUB	number of subintervals for partitioning the range of integration
SW	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
ALPHA	target significance level
MAXH	maximum number of interval halving steps to be carried out in finding the maximally raised nominal level

Details

The program uses 96-point Gauss-Legendre quadrature on each of the NSUB intervals into which the range of integration is partitioned.

Value

N	sample size
DEL0	noninferiority margin to the difference of the parameters of the marginal binomial distributions under comparison
K1	Parameter 1 of the Dirichlet prior for the family of trinomial distributions
K2	Parameter 2 of the Dirichlet prior for the family of trinomial distributions
K3	Parameter 3 of the Dirichlet prior for the family of trinomial distributions
NSUB	number of subintervals for partitioning the range of integration
SW	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
ALPHA	target significance level
MAXH	maximum number of interval halving steps to be carried out in finding the maximally raised nominal level
ALPHA0	result of the search for the largest admissible nominal level
SIZE0	size of the critical region corresponding to α_0
SIZE_UNC	size of the critical region of test at uncorrected nominal level α
POW	power against 7 different parameter configurations with $\delta = 0$

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>

Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 5.2.3.

Examples

```
mcnby_ni(25, .10, .5, .5, .5, 10, .05, .05, 5)
```

mcnby_ni_pp	<i>Computation of the posterior probability of the alternative hypothesis of noninferiority in the McNemar setting, given a specific point in the sample space</i>
-------------	--

Description

Evaluation of the integral on the right-hand side of Equation (5.24) on p. 88 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

```
mcnby_ni_pp(N, DEL0, N10, N01)
```

Arguments

N	sample size
DEL0	noninferiority margin to the difference of the parameters of the marginal binomial distributions under comparison
N10	count of pairs with $(X, Y) = (1, 0)$
N01	count of pairs with $(X, Y) = (0, 1)$

Details

The program uses 96-point Gauss-Legendre quadrature on each of 10 subintervals into which the range of integration is partitioned.

Value

N	sample size
DEL0	noninferiority margin to the difference of the parameters of the marginal binomial distributions under comparison
N10	count of pairs with $(X, Y) = (1, 0)$
N01	count of pairs with $(X, Y) = (0, 1)$
PPOST	posterior probability of the alternative hypothesis $K_1 : \delta > -\delta_0$ with respect to the noninformative prior determined according to Jeffrey's rule

Note

The program uses Equation (5.24) of Wellek S (2010) corrected for a typo in the middle line which must read

$$\int_{\delta_0}^{(1+\delta_0)/2} \left[B(n_{01} + 1/2, n - n_{01} + 1) p_{01}^{n_{01}-1/2} (1 - p_{01})^{n-n_{01}} \right]$$

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 5.2.3.

Examples

mcnby_ni_pp(72, 0.05, 4, 5)

mcnemasc

Determination of a corrected nominal significance level for the asymptotic test for equivalence of two paired binomial proportions with respect to the difference of their expectations (McNemar setting)

Description

The program computes the largest nominal significance level which can be substituted for the target level α without making the exact size of the asymptotic testing procedure larger than α .

Usage

mcnemasc(alpha, n, del0, sw, tol, maxh)

Arguments

alpha	significance level
n	sample size
del0	upper limit set to $ p_{10} - p_{01} $ under the alternative hypothesis of equivalence, with p_{10} and p_{01} denoting the probabilities of discordant pairs of both kinds
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
tol	upper bound to the absolute difference between size and target level below which the search for a corrected nominal level terminates
maxh	maximum number of interval halving steps to be carried out in finding the maximally raised nominal level

Value

alpha	significance level
n	sample size
de10	upper limit set to $ p_{10} - p_{01} $ under the alternative hypothesis of equivalence, with p_{10} and p_{01} denoting the probabilities of discordant pairs of both kinds
sw	width of the search grid for determining the maximum of the rejection probability on the common boundary of the hypotheses
ALPH_0	value of the corrected nominal level obtained after nh steps
NH	number of interval-halving steps actually performed
ERROR	error indicator messaging "!!!!!" if the sufficient condition for the correctness of the result output by the program was found violated

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 5.2.2.

Examples

```
mcnemasc(0.05, 50, 0.20, 0.05, 0.0005, 5)
```

mcnempow	<i>Exact rejection probability of the asymptotic test for equivalence of two paired binomial proportions with respect to the difference of their expectations (McNemar setting)</i>
----------	---

Description

The program computes exact values of the rejection probability of the asymptotic test for equivalence in the sense of $-\delta_0 < p_{10} - p_{01} < \delta_0$, at any nominal level α . [The largest α for which the test is valid in terms of the significance level, can be computed by means of the program mcnemasc.]

Usage

```
mcnempow(alpha, n, de10, p10, p01)
```

Arguments

alpha	nominal significance level
n	sample size
de10	upper limit set to $ \delta $ under the alternative hypothesis of equivalence
p10	true value of $P[X = 1, Y = 0]$
p01	true value of $P[X = 0, Y = 1]$

Value

alpha	nominal significance level
n	sample size
de10	upper limit set to $ \delta $ under the alternative hypothesis of equivalence
p10	true value of $P[X = 1, Y = 0]$
p01	true value of $P[X = 0, Y = 1]$
POW	exact rejection probability of the asymptotic McNemar test for equivalence at nominal level α
ERROR	error indicator messaging "!!!!!" if the sufficient condition for the correctness of the result output by the program was found violated

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>

Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, p.84.

Examples

```
mcnempow(0.024902, 50, 0.20, 0.30, 0.30)
```

mwtie_fr

Analogue of mwtie_xy for settings with grouped data

Description

Implementation of the asymptotically distribution-free test for equivalence of discrete distributions from which grouped data are obtained. Hypothesis formulation is in terms of the Mann-Whitney-Wilcoxon functional generalized to the case that ties between observations from different distributions may occur with positive probability. For details see Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition, p.155.

Usage

```
mwtie_fr(k,alpha,m,n,eps1_,eps2_,x,y)
```

Arguments

k	total number of grouped values which can be distinguished in the pooled sample
alpha	significance level
m	size of Sample 1
n	size of Sample 2
eps1_	absolute value of the left-hand limit of the hypothetical equivalence range for $\pi_+/(1 - \pi_0) - 1/2$
eps2_	right-hand limit of the hypothetical equivalence range for $\pi_+/(1 - \pi_0) - 1/2$
x	row vector with the m observations making up Sample1 as components
y	row vector with the n observations making up Sample2 as components

Details

Notation: π_+ and π_0 stands for the functional defined by $\pi_+ = P[X > Y]$ and $\pi_0 = P[X = Y]$, respectively, with $X \sim F \equiv$ cdf of Population 1 being independent of $Y \sim G \equiv$ cdf of Population 2.

Value

alpha	significance level
m	size of Sample 1
n	size of Sample 2
eps1_	absolute value of the left-hand limit of the hypothetical equivalence range for $\pi_+/(1 - \pi_0) - 1/2$
eps2_	right-hand limit of the hypothetical equivalence range for $\pi_+/(1 - \pi_0) - 1/2$
WXY_TIE	observed value of the U -statistics – based estimator of $\pi_+/(1 - \pi_0)$
SIGMAH	square root of the estimated asymptotic variance of $W_+/(1 - W_0)$
CRIT	upper critical bound to $ W_+/(1 - W_0) - 1/2 - (\varepsilon'_2 - \varepsilon'_1)/2 /\hat{\sigma}$
REJ	indicator of a positive [=1] vs negative [=0] rejection decision to be taken with the data under analysis

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

- Wellek S, Hampel B: A distribution-free two-sample equivalence test allowing for tied observations. *Biometrical Journal* 41 (1999), 171-186.
- Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.4.

Examples

```
x <- c(1,1,3,2,2,3,1,1,1,2,1,2,2,2,1,2,1,3,2,1,2,1,1,1,1,1,1,1,1,1,1,1,1,2,1,3,1,3,2,1,1,
      2,1,2,1,1,2,2,1,2,1,1,1,1,1,2,2,1,2,2,1,3,1,2,1,1,2,2,1,2,2,1,1,1,3,2,1,1,1,2,1,
      3,3,3,1,2,1,2,2,1,1,1,2,2,1,1,2,1,1,2,3,1,3,2,1,1,1,1,2,2,2,1,1,2,2,3,2,1,2,1,1,
      2,2,1,2,2,2,1,1,2,3,2,1,3,2,1,1,1,2,2,2,2,1,2,2,1,1,1,1,2,1,1,1,2,1,2,2,1,2,2,2,
      2,1,1,2,1,2,2,1,1,1,1,3,1,1,2,2,1,1,1,2,2,2,1,2,3,2,2,1,2,1,2,1,1,2,1,2,2,1,1,1,
      2,2,2,2)
y <- c(2,1,2,2,1,1,2,2,2,1,1,2,1,3,3,1,1,1,1,1,1,2,2,3,1,1,1,3,1,1,1,1,1,1,1,2,2,3,2,1,
      2,2,2,1,2,1,1,2,2,1,2,1,1,1,1,2,1,2,1,1,3,1,1,1,2,2,2,1,1,1,1,2,1,2,1,1,2,2,2,2,
      2,1,1,1,3,2,2,2,1,2,3,1,2,1,1,1,2,1,3,3,1,2,2,2,2,2,2,1,2,1,1,1,1,2,2,1,1,1,1,2,
      1,3,1,1,2,1,2,1,2,2,2,1,2,2,2,1,1,1,2,1,2,1,2,1,1,1,2,1,2,2,1,1,1,1,2,2,3,1,3,1,
      1,2,2,2,1,1,1,1,2,1,1,3,2,2,3,1,2,2,1,1,2,1,1,2,1,2,1,2,1,2,2,2,1,1,1,1,1,1,1,
      1,1,1,2,1,3,2,2,1,1,1,2,2,1,1,2,1,2,1,2,2,2,1,2,3,1,1,2,1,2,2,1,1,1,1,2,2,2,1,1,
      3,2,1,2,2,2,1,1,1,2,1,2,2,1,2,1,1,2)
mwtie_fr(3,0.05,204,258,0.10,0.10,x,y)
```

mwtie_xy

Distribution-free two-sample equivalence test for tied data: test statistic and critical upper bound

Description

Implementation of the asymptotically distribution-free test for equivalence of discrete distributions in terms of the Mann-Whitney-Wilcoxon functional generalized to the case that ties between observations from different distributions may occur with positive probability. For details see Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition, § 6.4.

Usage

```
mwtie_xy(alpha,m,n,eps1_,eps2_,x,y)
```

Arguments

alpha	significance level
m	size of Sample 1
n	size of Sample 2
eps1_	absolute value of the left-hand limit of the hypothetical equivalence range for $\pi_+/(1 - \pi_0) - 1/2$
eps2_	right-hand limit of the hypothetical equivalence range for $\pi_+/(1 - \pi_0) - 1/2$
x	row vector with the m observations making up Sample1 as components
y	row vector with the n observations making up Sample2 as components

Details

Notation: π_+ and π_0 stands for the functional defined by $\pi_+ = P[X > Y]$ and $\pi_0 = P[X = Y]$, respectively, with $X \sim F \equiv$ cdf of Population 1 being independent of $Y \sim G \equiv$ cdf of Population 2.

Value

alpha	significance level
m	size of Sample 1
n	size of Sample 2
eps1_	absolute value of the left-hand limit of the hypothetical equivalence range for $\pi_+/(1 - \pi_0) - 1/2$
eps2_	right-hand limit of the hypothetical equivalence range for $\pi_+/(1 - \pi_0) - 1/2$
WXY_TIE	observed value of the U -statistics – based estimator of $\pi_+/(1 - \pi_0)$
SIGMAH	square root of the estimated asymptotic variance of $W_+/(1 - W_0)$
CRIT	upper critical bound to $ W_+/(1 - W_0) - 1/2 - (\varepsilon'_2 - \varepsilon'_1)/2 /\hat{\sigma}$
REJ	indicator of a positive [=1] vs negative [=0] rejection decision to be taken with the data under analysis

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S, Hampel B: A distribution-free two-sample equivalence test allowing for tied observations. *Biometrical Journal* 41 (1999), 171-186.

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.4.

Examples

```
x <- c(1,1,3,2,2,3,1,1,1,2)
y <- c(2,1,2,2,1,1,2,2,2,1,1,2)
mwtie_xy(0.05,10,12,0.10,0.10,x,y)
```

postmys	<i>Bayesian posterior probability of the alternative hypothesis in the setting of the one-sample t-test for equivalence</i>
---------	---

Description

Evaluation of the integral appearing on the right-hand side of equation (3.6) on p. 38 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition

Usage

```
postmys(n,dq,sd,eps1,eps2,tol)
```

Arguments

n	sample size
dq	mean within-pair difference observed in the sample under analysis
sd	square root of the sample variance of the within-pair differences
eps1	absolute value of the left-hand limit of the hypothetical equivalence range for δ/σ_D
eps2	right-hand limit of the hypothetical equivalence range for δ/σ_D
tol	tolerance for the error induced through truncating the range of integration on the right

Details

The program uses 96-point Gauss-Legendre quadrature.

Value

n	sample size
dq	mean within-pair difference observed in the sample under analysis
sd	square root of the sample variance of the within-pair differences
eps1	absolute value of the left-hand limit of the hypothetical equivalence range for δ/σ_D
eps2	right-hand limit of the hypothetical equivalence range for δ/σ_D
tol	tolerance for the error induced through truncating the range of integration on the right
PPOST	posterior probability of the set of all (δ, σ_D) such that $-\varepsilon_1 < \delta/\sigma_D < \varepsilon_2$

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 3.2.

Examples

```
postmys(23, 0.16, 3.99, 0.5, 0.5, 1e-6)
```

powsign	<i>Nonconditional power of the UMPU sign test for equivalence and its nonrandomized counterpart</i>
---------	---

Description

The program computes for each possible value of the number n_0 of zero observations the power conditional on $N_0 = n_0$ and averages these conditional power values with respect to the distribution of N_0 . Equivalence is defined in terms of the logarithm of the ratio p_+/p_- , where p_+ and p_- denotes the probability of obtaining a positive and negative sign, respectively.

Usage

```
powsign(alpha,n,eps1,eps2,poa)
```

Arguments

alpha	significance level
n	sample size
eps1	absolute value of the lower limit of the hypothetical equivalence range for $\log(p_+/p_-)$.
eps2	upper limit of the hypothetical equivalence range for $\log(p_+/p_-)$.
poa	probability of a tie under the alternative of interest

Value

alpha	significance level
n	sample size
eps1	absolute value of the lower limit of the hypothetical equivalence range for $\log(p_+/p_-)$.
eps2	upper limit of the hypothetical equivalence range for $\log(p_+/p_-)$.
poa	probability of a tie under the alternative of interest
POWNONRD	power of the nonrandomized version of the test against the alternative $p_+ = p_- = (1 - p_0)/2$
POW	power of the randomized UMPU test against the alternative $p_+ = p_- = (1 - p_0)/2$

Note

A special case of the test whose power is computed by this program, is the exact conditional equivalence test for the McNemar setting (cf. Wellek 2010, pp. 76-77).

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 5.1.

Examples

```
powsign(0.06580, 50, 0.847298, 0.847298, 0.26)
```

pow_abe	<i>Confidence interval inclusion test for average bioequivalence: exact power against an arbitrary specific alternative</i>
---------	---

Description

Evaluation of the integral on the right-hand side of equation (10.11) of p. 317 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition

Usage

```
pow_abe(m, n, alpha, del_0, del, sig)
```

Arguments

m	sample size in sequence group T(est)/R(eference)
n	sample size in sequence group R(eference)/T(est)
alpha	significance level
del_0	equivalence margin to the absolute value of the log-ratio μ_T^* and μ_R^* of the formulation effects
del	assumed true value of $ \log(\mu_T^*/\mu_R^*) $, with $0 \leq \delta < \delta_0$
sig	theoretical standard deviation of the log within-subject bioavailability ratios in each sequence group

Details

The program uses 96-point Gauss-Legendre quadrature.

Value

m	sample size in sequence group T(est)/R(eference)
n	sample size in sequence group R(eference)/T(est)
alpha	significance level
del_0	equivalence margin to the absolute value of the log-ratio μ_T^* and μ_R^* of the formulation effects
del	assumed true value of $ \log(\mu_T^*/\mu_R^*) $, with $0 \leq \delta < \delta_0$
POW_ABE	power of the interval inclusion test for average bioequivalence against the specific alternative given by (δ, σ)

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 10.2.1.

Examples

```
pow_abe(12, 13, 0.05, log(1.25), log(1.25)/2, 0.175624)
```

po_pbibe	<i>Bayesian posterior probability of the alternative hypothesis of probability-based individual bioequivalence (PBIBE)</i>
----------	--

Description

Implementation of the algorithm presented in § 10.3.3 of Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition.

Usage

```
po_pbibe(n, eps, pio, zq, s, tol, sw, ihmax)
```

Arguments

n	sample size
eps	equivalence margin to an individual log-bioavailability ratio
pio	prespecified lower bound to the probability of obtaining an individual log-bioavailability ratio falling in the equivalence range $(-\varepsilon, \varepsilon)$
zq	mean log-bioavailability ratio observed in the sample under analysis
s	square root of the sample variance of the log-bioavailability ratios
tol	maximum numerical error allowed for transforming the hypothesis of PBIBE into a region in the parameter space of the log-normal distribution assumed to underlie the given sample of individual bioavailability ratios
sw	step width used in the numerical procedure yielding results at a level of accuracy specified by the value chosen for tol
ihmax	maximum number of interval halving steps to be carried out in finding the region specified in the parameter space according to the criterion of PBIBE

Details

The program uses 96-point Gauss-Legendre quadrature.

Value

n	sample size
eps	equivalence margin to an individual log-bioavailability ratio
pio	prespecified lower bound to the probability of obtaining an individual log-bioavailability ratio falling in the equivalence range $(-\varepsilon, \varepsilon)$
zq	mean log-bioavailability ratio observed in the sample under analysis
s	square root of the sample variance of the log-bioavailability ratios
tol	maximum numerical error allowed for transforming the hypothesis of PBIBE into a region in the parameter space of the log-normal distribution assumed to underlie the given sample of individual bioavailability ratios
sw	step width used in the numerical procedure yielding results at a level of accuracy specified by the value chosen for tol
ihmax	maximum number of interval halving steps to be carried out in finding the region specified in the parameter space according to the criterion of PBIBE
PO_PBIBE	posterior probability of the alternative hypothesis of PBIBE

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Bayesian construction of an improved parametric test for probability-based individual bioequivalence. *Biometrical Journal* 42 (2000), 1039-52.

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 10.3.3.

Examples

```
po_pbibe(20,0.25,0.75,0.17451,0.04169, 10e-10,0.01,100)
```

sgnrk	<i>Signed rank test for equivalence of an arbitrary continuous distribution of the intraindividual differences in terms of the probability of a positive sign of a Walsh average: test statistic and critical upper bound</i>
-------	---

Description

Implementation of the paired-data analogue of the Mann-Whitney-Wilcoxon test for equivalence of continuous distributions. The continuity assumption relates to the intraindividual differences D_i . For details see Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition, § 5.4.

Usage

```
sgnrk(alpha,n,qp1,qp12,d)
```

Arguments

alpha	significance level
n	sample size
qp1	lower equivalence limit q'_+ to the target functional q_+
qp12	upper equivalence limit q''_+ to the target functional q_+
d	row vector with the intraindividual differences for all n pairs as components

Details

q_+ is the probability of getting a positive sign of the so-called Walsh-average of a pair of within-subject differences and can be viewed as a natural paired-observations analogue of the Mann-Whitney functional $\pi_+ = P[X > Y]$.

Value

alpha	significance level
n	sample size
qp1	lower equivalence limit q'_+ to the target functional q_+
qp12	upper equivalence limit q''_+ to the target functional q_+
U_p1	observed value of the U -statistics estimator of q_+
SIGMAH	square root of the estimated asymptotic variance of U_+
CRIT	upper critical bound to $ U_+ - (q'_+ + q''_+)/2 /\hat{\sigma}$
REJ	indicator of a positive [=1] vs negative [=0] rejection decision to be taken with the data under analysis

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 5.4.

Examples

```
d <- c(-0.5,0.333,0.667,1.333,1.5,-2.0,-1.0,-0.167,1.667,0.833,-2.167,-1.833,
      4.5,-7.5,2.667,3.333,-4.167,5.667,2.333,-2.5)
sgnrk(0.05,20,0.2398,0.7602,d)
```

srktie_d	<i>Generalized signed rank test for equivalence for tied data: test statistic and critical upper bound</i>
----------	--

Description

Implementation of a generalized version of the signed-rank test for equivalence allowing for arbitrary patterns of ties between the within-subject differences. For details see Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition, § 5.5.

Usage

```
srktie_d(n,alpha,eps1,eps2,d)
```

Arguments

n	sample size
alpha	significance level
eps1	absolute value of the left-hand limit of the hypothetical equivalence range for $q_+/(1 - q_0) - 1/2$
eps2	right-hand limit of the hypothetical equivalence range for $q_+/(1 - q_0) - 1/2$
d	row vector with the intraindividual differences for all n pairs as components

Details

Notation: q_+ and q_0 stands for the functional defined by $q_+ = P[D_i + D_j > 0]$ and $q_0 = P[D_i + D_j = 0]$, respectively, with D_i and D_j as the intraindividual differences observed in two individuals independently selected from the underlying bivariate population.

Value

n	sample size
alpha	significance level
eps1	absolute value of the left-hand limit of the hypothetical equivalence range for $q_+/(1 - q_0) - 1/2$
eps2	right-hand limit of the hypothetical equivalence range for $q_+/(1 - q_0) - 1/2$
U_p1	observed value of the U -statistics estimator of q_+
U_0	observed value of the U -statistics estimator of q_0
UAS_PL	observed value of $U_+/(1 - U_0)$
TAUHAS	square root of the estimated asymptotic variance of $\sqrt{n}U_+/(1 - U_0)$
CRIT	upper critical bound to $\sqrt{n} U_+/(1 - U_0) - 1/2 - (\varepsilon_2 - \varepsilon_1)/2 /\hat{\tau}$
REJ	indicator of a positive [=1] vs negative [=0] rejection decision to be taken with the data under analysis

Note

The function `srktie_d` can be viewed as the paired-data analogue of `mwtie_xy`

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 5.5.

Examples

```
d <- c(0.8, 0.2, 0.0, -0.1, -0.3, 0.3, -0.1, 0.4, 0.6, 0.2, 0.0, -0.2, -0.3, 0.0, 0.1, 0.3, -0.3,
      0.1, -0.2, -0.5, 0.2, -0.1, 0.2, -0.1)
srktie_d(24, 0.05, 0.2602, 0.2602, d)
```

srktie_m

Analogue of srktie_d for settings where the distribution of intraindividual differences is concentrated on a finite lattice

Description

Analogue of the function `srktie_d` tailored for settings where the distribution of the within-subject differences is concentrated on a finite lattice. For details see Wellek S (2010) Testing statistical hypotheses of equivalence and noninferiority. Second edition, pp.112-3.

Usage

```
srktie_m(n, alpha, eps1, eps2, w, d)
```

Arguments

n	sample size
alpha	significance level
eps1	absolute value of the left-hand limit of the hypothetical equivalence range for $q_+/(1 - q_0) - 1/2$
eps2	right-hand limit of the hypothetical equivalence range for $q_+/(1 - q_0) - 1/2$
w	span of the lattice in which the intraindividual differences take their values
d	row vector with the intraindividual differences for all n pairs as components

Details

Notation: q_+ and q_0 stands for the functional defined by $q_+ = P[D_i + D_j > 0]$ and $q_0 = P[D_i + D_j = 0]$, respectively, with D_i and D_j as the intraindividual differences observed in two individuals independently selected from the underlying bivariate population.

Value

n	sample size
alpha	significance level
eps1	absolute value of the left-hand limit of the hypothetical equivalence range for $q_+/(1 - q_0) - 1/2$
eps2	right-hand limit of the hypothetical equivalence range for $q_+/(1 - q_0) - 1/2$
w	span of the lattice in which the intraindividual differences take their values
U_p1	observed value of the U -statistics estimator of q_+
U_0	observed value of the U -statistics estimator of q_0
UAS_PL	observed value of $U_+/(1 - U_0)$
TAUHAS	square root of the estimated asymptotic variance of $\sqrt{n}U_+/(1 - U_0)$
CRIT	upper critical bound to $\sqrt{n} U_+/(1 - U_0) - 1/2 - (\varepsilon_2 - \varepsilon_1)/2 /\hat{\tau}$
REJ	indicator of a positive [=1] vs negative [=0] rejection decision to be taken with the data under analysis

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, pp. 112-114.

Examples

```
d <- c(0.8,0.2,0.0,-0.1,-0.3,0.3,-0.1,0.4,0.6,0.2,0.0,-0.2,-0.3,0.0,0.1,0.3,-0.3,
      0.1,-0.2,-0.5,0.2,-0.1,0.2,-0.1)
srktie_m(24,0.05,0.2602,0.2602,0.1,d)
```

tt1st	<i>Critical constants and power against the null alternative of the one-sample t-test for equivalence with an arbitrary, maybe nonsymmetric choice of the limits of the equivalence range</i>
-------	--

Description

The function computes the critical constants defining the uniformly most powerful invariant test for the problem $\delta/\sigma_D \leq \theta_1$ or $\delta/\sigma_D \geq \theta_2$ versus $\theta_1 < \delta/\sigma_D < \theta_2$, with (θ_1, θ_2) as a fixed nondegenerate interval on the real line. In addition, tt1st outputs the power against the null alternative $\delta = 0$.

Usage

```
tt1st(n,alpha,theta1,theta2,tol,itmax)
```

Arguments

n	sample size
alpha	significance level
theta1	lower equivalence limit to δ/σ_D
theta2	upper equivalence limit to δ/σ_D
tol	tolerable deviation from α of the rejection probability at either boundary of the hypothetical equivalence interval
itmax	maximum number of iteration steps

Value

n	sample size
alpha	significance level
theta1	lower equivalence limit to δ/σ_D
theta2	upper equivalence limit to δ/σ_D
IT	number of iteration steps performed until reaching the stopping criterion corresponding to TOL
C1	left-hand limit of the critical interval for the one-sample t -statistic
C2	right-hand limit of the critical interval for the one-sample t -statistic
ERR1	deviation of the rejection probability from α under $\delta/\sigma_D = \theta_1$
ERR2	deviation of the rejection probability from α under $\delta/\sigma_D = \theta_2$
POW0	power of the UMPI test against the alternative $\delta = 0$

Note

If the output value of ERR2 is NA, the deviation of the rejection probability at the right-hand boundary of the hypothetical equivalence interval from α is smaller than the smallest real number representable in R.

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 5.3.

Examples

```
tt1st(36,0.05, -0.4716,0.3853,1e-10,50)
```

 tt2st

Critical constants and power against the null alternative of the two-sample t-test for equivalence with an arbitrary, maybe nonsymmetric choice of the limits of the equivalence range

Description

The function computes the critical constants defining the uniformly most powerful invariant test for the problem $(\xi - \eta)/\sigma \leq -\varepsilon_1$ or $(\xi - \eta)/\sigma \geq \varepsilon_2$ versus $-\varepsilon_1 < (\xi - \eta)/\sigma < \varepsilon_2$, with ξ and η denoting the expected values of two normal distributions with common variance σ^2 from which independent samples are taken. In addition, tt2st outputs the power against the null alternative $\xi = \eta$.

Usage

```
tt2st(m,n,alpha,eps1,eps2,tol,itmax)
```

Arguments

m	size of the sample from $\mathcal{N}(\xi, \sigma^2)$
n	size of the sample from $\mathcal{N}(\eta, \sigma^2)$
alpha	significance level
eps1	absolute value of the lower equivalence limit to $(\xi - \eta)/\sigma$
eps2	upper equivalence limit to $(\xi - \eta)/\sigma$
tol	tolerable deviation from α of the rejection probability at either boundary of the hypothetical equivalence interval
itmax	maximum number of iteration steps

Value

m	size of the sample from $\mathcal{N}(\xi, \sigma^2)$
n	size of the sample from $\mathcal{N}(\eta, \sigma^2)$
alpha	significance level
eps1	absolute value of the lower equivalence limit to $(\xi - \eta)/\sigma$
eps2	upper equivalence limit to $(\xi - \eta)/\sigma$
IT	number of iteration steps performed until reaching the stopping criterion corresponding to TOL
C1	left-hand limit of the critical interval for the two-sample t -statistic
C2	right-hand limit of the critical interval for the two-sample t -statistic
ERR1	deviation of the rejection probability from α under $(\xi - \eta)/\sigma = -\varepsilon_1$
ERR2	deviation of the rejection probability from α under $(\xi - \eta)/\sigma = \varepsilon_2$
POW0	power of the UMPI test against the alternative $\xi = \eta$

Note

If the output value of ERR2 is NA, the deviation of the rejection probability at the right-hand boundary of the hypothetical equivalence interval from α is smaller than the smallest real number representable in R.

Author(s)

Stefan Wellek <stefan.wellek@zi-mannheim.de>
 Peter Ziegler <peter.ziegler@zi-mannheim.de>

References

Wellek S: Testing statistical hypotheses of equivalence and noninferiority. Second edition. Boca Raton: Chapman & Hall/CRC Press, 2010, § 6.1.

Examples

```
tt2st(12, 12, 0.05, 0.50, 1.00, 1e-10, 50)
```

Index

- * **2x2 contingency table**
 - gofind_t, 30
- * **Bayesian test**
 - bi2by_ni_del, 8
 - bi2by_ni_OR, 10
 - mcnby_ni, 35
 - po_pbibe, 47
 - postmys, 43
- * **Dirichlet prior**
 - mcnby_ni, 35
 - mcnby_ni_pp, 37
- * **Euclidean distance statistic**
 - gofind_t, 30
 - gofsimpt, 31
- * **Gauss-Legendre quadrature**
 - bi2by_ni_del, 8
 - bi2by_ni_OR, 10
 - mcnby_ni, 35
 - mcnby_ni_pp, 37
 - po_pbibe, 47
 - postmys, 43
- * **Gaussian data**
 - fstretch, 26
- * **Hardy-Weinberg equilibrium (HWE)**
 - gofhwex, 28
 - gofhwex_1s, 29
- * **Hardy-Weinberg equilibrium**
 - cf_reh_exact, 22
 - cf_reh_midp, 23
- * **Jeffrey's rule**
 - mcnby_ni_pp, 37
- * **Jeffreys prior**
 - postmys, 43
- * **Mann-Whitney functional**
 - mawi, 33
 - mwtie_fr, 40
 - mwtie_xy, 42
- * **McNemar setting**
 - mcnasc_ni, 34
 - mcnby_ni_pp, 37
 - mcnemasc, 38
 - mcnempow, 39
 - powsign, 45
- * **U-statistics estimators**
 - mwtie_fr, 40
 - mwtie_xy, 42
 - srktie_d, 50
 - srktie_m, 51
- * **UMP test for equivalence**
 - exp1st, 25
- * **UMP test**
 - bi1st, 3
- * **UMPU test**
 - bi2st, 16
 - powsign, 45
- * **Wald-type**
 - bi2wld_ni_del, 21
- * **Walsh averages**
 - sgnrk, 48
 - srktie_d, 50
 - srktie_m, 51
- * **asymptotic normality**
 - gofind_t, 30
 - gofsimpt, 31
 - mawi, 33
 - mwtie_fr, 40
 - mwtie_xy, 42
 - sgnrk, 48
 - srktie_d, 50
 - srktie_m, 51
- * **asymptotic test for equivalence**
 - mcnempow, 39
- * **asymptotic test**
 - bi2diffac, 11
 - bi2dipow, 12
 - mcnasc_ni, 34
 - mcnemasc, 38
- * **average bioequivalence**

- pow_abe, 46
- * **binary observations**
 - mcnasc_ni, 34
 - mcnemasc, 38
 - powsign, 45
- * **binomial one-sample problem**
 - bi1st, 3
- * **binomial two-sample problem**
 - bi2aeq1, 4
 - bi2aeq2, 5
 - bi2aeq3, 7
 - bi2diffac, 11
 - bi2dipow, 12
 - bi2rlv1, 14
 - bi2rlv2, 15
 - bi2st, 16
 - bi2ste1, 17
 - bi2ste2, 18
 - bi2ste3, 19
- * **conditional test**
 - bi2st, 16
- * **confidence interval inclusion test**
 - pow_abe, 46
- * **confidence interval inclusion**
 - cf_reh_exact, 22
 - cf_reh_midp, 23
- * **continuous observations**
 - mawi, 33
 - sgnrk, 48
- * **corrected nominal level**
 - bi2diffac, 11
 - mcnasc_ni, 34
 - mcnemasc, 38
- * **difference between response probabilities**
 - mcnby_ni_pp, 37
- * **difference of proportions**
 - bi2by_ni_del, 8
 - mcnby_ni, 35
- * **difference of success probabilities**
 - bi2diffac, 11
 - bi2dipow, 12
- * **dispersion equivalence**
 - fstretch, 26
- * **equivalence test**
 - cf_reh_exact, 22
 - cf_reh_midp, 23
- * **equivalence**
 - bi1st, 3
 - bi2aeq1, 4
 - bi2aeq2, 5
 - bi2aeq3, 7
 - bi2diffac, 11
 - bi2dipow, 12
 - bi2st, 16
 - mcnemasc, 38
 - powsign, 45
- * **establishing approximate independence of two categorical variables**
 - gofind_t, 30
- * **exact Fisher-type test nonrandomized version**
 - bi2aeq3, 7
 - bi2ste3, 19
- * **exact Fisher-type test**
 - bi2aeq1, 4
 - bi2aeq2, 5
 - bi2rlv1, 14
 - bi2rlv2, 15
 - bi2st, 16
 - bi2ste1, 17
 - bi2ste2, 18
- * **exact conditional confidence limit**
 - cf_reh_exact, 22
- * **exact conditional test**
 - powsign, 45
- * **exact power**
 - bi2dipow, 12
 - mcnempow, 39
- * **exponential distribution**
 - exp1st, 25
- * **finite lattice**
 - srktie_m, 51
- * **fully specified multinomial distribution**
 - gofsimpt, 31
- * **genetic association study**
 - cf_reh_exact, 22
 - cf_reh_midp, 23
- * **goodness of fit**
 - cf_reh_exact, 22
 - cf_reh_midp, 23
- * **homoskedasticity**
 - tt2st, 54
- * **individual bioequivalence**
 - po_pbibe, 47
- * **lognormal distribution**
 - po_pbibe, 47

- pow_abe, 46
- * **matched-pair design with binary data**
 - mcnby_ni, 35
- * **maximally raised nominal level**
 - bi2aeq3, 7
 - bi2ste3, 19
- * **mid-p value**
 - cf_reh_midp, 23
- * **model validation**
 - gofhwex, 28
- * **model verification**
 - gofsimpt, 31
- * **nominal level**
 - bi2wld_ni_del, 21
- * **noncentral t-distribution**
 - tt1st, 53
 - tt2st, 54
- * **noninferiority**
 - bi2by_ni_del, 8
 - bi2by_ni_OR, 10
 - bi2ste1, 17
 - bi2ste2, 18
 - bi2ste3, 19
 - gofhwex_1s, 29
 - mcnasc_ni, 34
 - mcnby_ni, 35
 - mcnby_ni_pp, 37
- * **nonparametric paired-sample equivalence test**
 - sgnrk, 48
 - srktie_d, 50
 - srktie_m, 51
- * **nonparametric two-sample equivalence test**
 - mawi, 33
 - mwtie_fr, 40
 - mwtie_xy, 42
- * **normal distribution**
 - postmys, 43
- * **odds ratio**
 - bi2aeq1, 4
 - bi2aeq2, 5
 - bi2aeq3, 7
 - bi2by_ni_OR, 10
 - bi2rlv1, 14
 - bi2rlv2, 15
 - bi2st, 16
 - bi2ste1, 17
 - bi2ste2, 18
 - bi2ste3, 19
- * **one-dimensional contingency table**
 - gofsimpt, 31
- * **one-sample problem**
 - exp1st, 25
- * **package**
 - EQUIVNONINF-package, 2
- * **paired data**
 - postmys, 43
- * **paired samples**
 - mcnasc_ni, 34
 - mcnemasc, 38
 - powsign, 45
- * **paired t-test for equivalence**
 - tt1st, 53
- * **parallel-group design with binary data**
 - bi2by_ni_del, 8
 - bi2by_ni_OR, 10
- * **posterior probability of the alternative hypothesis**
 - mcnby_ni_pp, 37
- * **power**
 - bi1st, 3
 - bi2aeq1, 4
 - bi2rlv1, 14
 - bi2ste1, 17
 - exp1st, 25
 - fstretch, 26
 - tt1st, 53
 - tt2st, 54
- * **product of Jeffrey's priors**
 - bi2by_ni_del, 8
 - bi2by_ni_OR, 10
- * **randomized test**
 - bi1st, 3
 - bi2st, 16
- * **relative excess heterozygosity (REH)**
 - gofhwex_1s, 29
- * **relative excess heterozygosity**
 - cf_reh_exact, 22
 - cf_reh_midp, 23
- * **relevant differences**
 - bi2rlv1, 14
 - bi2rlv2, 15
- * **sample size**
 - bi2aeq2, 5
 - bi2rlv2, 15

- bi2ste2, 18
 - * **single nucleotide polymorphism (SNP)**
 - gofhwex, 28
 - gofhwex_1s, 29
 - * **standardized difference of means**
 - postmys, 43
 - tt1st, 53
 - tt2st, 54
 - * **tied observations**
 - mwtie_fr, 40
 - mwtie_xy, 42
 - srktie_d, 50
 - srktie_m, 51
 - * **two-period crossover**
 - pow_abe, 46
 - * **two-sample problem**
 - fstretch, 26
 - * **two-sample t-test for equivalence**
 - tt2st, 54
 - * **uniformly most powerful invariant test**
 - fstretch, 26
 - tt1st, 53
 - tt2st, 54
 - * **uniformly most powerful unbiased test**
 - gofhwex, 28
 - gofhwex_1s, 29
-
- bi1st, 3
 - bi2aeq1, 4
 - bi2aeq2, 5
 - bi2aeq3, 7
 - bi2by_ni_del, 8
 - bi2by_ni_OR, 10
 - bi2diffac, 11
 - bi2dipow, 12
 - bi2rlv1, 14
 - bi2rlv2, 15
 - bi2st, 16
 - bi2ste1, 17
 - bi2ste2, 18
 - bi2ste3, 19
 - bi2wld_ni_del, 21
-
- cf_reh_exact, 22
 - cf_reh_midp, 23
-
- EQUIVNONINF (EQUIVNONINF-package), 2
 - EQUIVNONINF-package, 2
 - exp1st, 25
 - fstretch, 26
 - gofhwex, 28
 - gofhwex_1s, 29
 - gofind_t, 30
 - gofsimpt, 31
 - mawi, 33
 - mcnasc_ni, 34
 - mcnby_ni, 35
 - mcnby_ni_pp, 37
 - mcnemasc, 38
 - mcnempow, 39
 - mwtie_fr, 40
 - mwtie_xy, 42
 - po_pbibe, 47
 - postmys, 43
 - pow_abe, 46
 - powsign, 45
 - sgnrk, 48
 - srktie_d, 50
 - srktie_m, 51
 - tt1st, 53
 - tt2st, 54