

# Package: SPIChanges (via r-universe)

March 7, 2025

**Title** Improves the Interpretation of the Standardized Precipitation Index Under Changing Climate Conditions

**Version** 0.1.0

**Description** Improves the interpretation of the Standardized Precipitation Index under changing climate conditions. The package uses the nonstationary approach proposed in Blain et al. (2022) <[doi:10.1002/joc.7550](https://doi.org/10.1002/joc.7550)> to detect trends in rainfall quantities and to quantify the effect of such trends on the probability of a drought event occurring.

**License** MIT + file LICENSE

**URL** <https://github.com/gabrielblain/SPIChanges>

**BugReports** <https://github.com/gabrielblain/SPIChanges/issues>

**Depends** R (>= 2.10)

**Imports** gamlss, gamlss.dist, lubridate, MuMIn, rlang, spsUtil, stats, brglm2, zoo

**Suggests** archive, curl, doParallel, dplyr, foreach, ggplot2, grid, knitr, ncdf4, patchwork, RColorBrewer, rmarkdown, roxygen2, sf, purrr, testthat (>= 3.0.0), tidy

**VignetteBuilder** knitr

**Config/roxygen2/filename** globals.R

**Config/roxygen2/unique** FALSE

**Config/testthat/edition** 3

**Encoding** UTF-8

**Language** en-US

**LazyData** true

**RoxygenNote** 7.3.2

**NeedsCompilation** no

**Author** Gabriel Constantino Blain [aut, cre]  
(<<https://orcid.org/0000-0001-8832-7734>>), Graciela R. Sobierajski [aut] (<<https://orcid.org/0000-0002-7211-9268>>),

Leticia L. Martins [aut]  
 (<<https://orcid.org/0000-0002-0299-3005>>), Adam H. Sparks [aut]  
 (<<https://orcid.org/0000-0002-0061-8359>>)

**Maintainer** Gabriel Constantino Blain <[gabriel.blain@sp.gov.br](mailto:gabriel.blain@sp.gov.br)>

**Repository** CRAN

**Date/Publication** 2025-02-05 18:10:08 UTC

**Config/pak/sysreqs** libicu-dev libssl-dev

## Contents

|                        |          |
|------------------------|----------|
| CampinasRain . . . . . | 2        |
| lonlat . . . . .       | 3        |
| Map . . . . .          | 3        |
| rainTS4 . . . . .      | 4        |
| SPIChanges . . . . .   | 5        |
| TSaggreg . . . . .     | 6        |
| <b>Index</b>           | <b>7</b> |

---

|              |                               |
|--------------|-------------------------------|
| CampinasRain | <i>Daily Rainfall Amounts</i> |
|--------------|-------------------------------|

---

## Description

Daily rainfall amounts recorded in millimetre in Campinas state of Sao Paulo, Brazil.

## Usage

CampinasRain

## Format

CampinasRain:  
 A dataframe with 2 columns and 16071 rows:

**Date** Date

**Rain** Daily rainfall amounts recorded in millimetre in Campinas state of Sao Paulo, Brazil ...

## Source

<https://psl.noaa.gov/>

---

|        |                                      |
|--------|--------------------------------------|
| lonlat | <i>Coordinates for Entire Brazil</i> |
|--------|--------------------------------------|

---

**Description**

This dataset provides coordinates for Brazil at a spatial resolution of 0.5 x 0.5 degrees.

**Usage**

lonlat

**Format**

lonlat:

A matrix with 2841 rows and 2 columns:

**lon** Longitude in decimal degrees

**lat** Latitude in decimal degrees ...

**Source**

Data generated using information from the Instituto Brasileiro de Geografia e Estatística (IBGE). For more information, visit: <https://www.ibge.gov.br/>

---

|     |  |
|-----|--|
| Map | <i>Changes in Drought Events in Brazil (1980-2024)</i> |
|-----|--|

---

**Description**

This dataset provides the changes (in percentage) in Moderate to Extreme, Severe to Extreme, and Extreme Drought events across Brazil for the four seasons (Summer, Autumn, Winter, and Spring) over the period 1980–2024. The data were generated using the SPIChanges() function applied to rainfall data from NOAA PSL.

**Usage**

Map

**Format**

Map:

A matrix with 2841 rows and 18 columns:

**lon** Longitude in decimal degrees

**lat** Latitude in decimal degrees

**SummerModerate** Percentage change in moderate drought events in Summer

**SummerSevere** Percentage change in severe drought events in Summer  
**SummerExtreme** Percentage change in extreme drought events in Summer  
**SummerModel** The selected gamma-based model in Summer  
**AutumnModerate** Percentage change in moderate drought events in Autumn  
**AutumnSevere** Percentage change in severe drought events in Autumn  
**AutumnExtreme** Percentage change in extreme drought events in Autumn  
**AutumnModel** The selected gamma-based model in Autumn  
**WinterModerate** Percentage change in moderate drought events in Winter  
**WinterSevere** Percentage change in severe drought events in Winter  
**WinterExtreme** Percentage change in extreme drought events in Winter  
**WinterModel** The selected gamma-based model in Winter  
**SpringModerate** Percentage change in moderate drought events in Spring  
**SpringSevere** Percentage change in severe drought events in Spring  
**SpringExtreme** Percentage change in extreme drought events in Spring  
**SpringModel** The selected gamma-based model in Spring ...

#### Source

Generated using the SPIChanges() function applied to rainfall data obtained from NOAA PSL (<https://psl.noaa.gov/>).

---

rainTS4

*Rainfall Aggregated at the 4-quasi week time scale*

---

#### Description

Rainfall amounts aggregated using TSaggreg(). Campinas state of Sao Paulo, Brazil.

#### Usage

```
rainTS4
```

#### Format

**rainTS4:**  
 A matrix with 4 columns and 576 rows:  
**Year** Year  
**Month** Month  
**quasiWeek** Four quasi-weekly periods in each month  
**rain.at.TS4** Rainfall amounts aggregated at TS equal to four ...

#### Source

<https://psl.noaa.gov/>

---

|            |   |
|------------|---|
| SPIChanges | <i>Detect trends and quantify their effect on the probability of SPI values occurring</i> |
|------------|---|

---

### Description

Detect trends and quantify their effect on the probability of SPI values occurring

### Usage

```
SPIChanges(rain.at.TS, only.linear = "Yes")
```

### Arguments

|             |   |
|-------------|---|
| rain.at.TS  | A 4-column matrix generated with TSaggreg(). No other objects are accepted. <ul style="list-style-type: none"> <li>• 1st column is years (YYYY),</li> <li>• 2nd is the months (1 to 12),</li> <li>• 3rd is the quasiWeeks (1 to 4),</li> <li>• and 4th is the rainfall totals accumulated at a time scale.</li> </ul> |
| only.linear | A character string value (Yes or No) defining if the function must consider only linear models (Yes) or linear and non-linear models (No). Default is Yes.  |

### Value

A list object with:

**data.week** The Rainfall amounts, SPI, cumulative probability of the SPI values under the stationary approach, cumulative probability of the SPI values under the non-stationary approach, and the changes in the frequency of below zero SPI values caused by the changes in rainfall patterns.

**model.selection** The generalized additive model that best fits the rainfall series

**Changes.Freq.Drought** changes in the frequency of zero precipitation, moderate to extreme, severe to extreme and extreme drought events, as categorized by the SPI classification system, caused by the changes in rainfall patterns. Changes in the precipitation amounts associated describing normal conditions is also shown.

**Statistics** Year to year changes in the expected frequency of moderate to extreme, severe to extreme and extreme drought events.

**data.week** The Rainfall amounts, SPI, cumulative probability of the SPI values under the stationary approach, cumulative probability of the SPI values under the non-stationary approach, and the changes in the frequency of below zero SPI values caused by the changes in rainfall patterns.

**model.selection** The generalized additive model that best fits the rainfall series

**Changes.Freq.Drought** changes in the frequency of zero precipitation, moderate, severe and extreme drought events, as defined by the SPI classification system, caused by the changes in rainfall patterns. Changes in the precipitation amounts associated describing normal conditions is also shown.

**Statistics** Year to year changes in the expected frequency of moderate, severe and extreme drought events.

**Examples**

```
rainTS4 <- rainTS4
Changes_SPI <- SPIChanges(rain.at.TS=rainTS4, only.linear = "yes")
```

---

**TSaggreg***Aggregates daily rainfall totals at quasi-week time scales*

---

**Description**

Aggregates daily rainfall totals at quasi-week time scales

**Usage**

```
TSaggreg(daily.rain, start.date, TS = 4L)
```

**Arguments**

|                         |  |
|-------------------------|--|
| <code>daily.rain</code> | Vector, 1-column matrix or data frame with daily rainfall totals.  |
| <code>start.date</code> | Date at which the aggregation should start. Preferred formats are “YYYY-MM-DD”, “YYYY/MM/DD” but most any valid date format should work. |
| <code>TS</code>         | Time scale on the quasiWeek basis (integer values between 1 and 96). Default is 4, which corresponds to the monthly time scale.          |

**Value**

A matrix with rainfall amounts aggregated at the time scale selected by the user

**Examples**

```
daily.rain <- CampinasRain[,2]
rainTS4 <- TSaggreg(daily.rain=daily.rain, start.date="1980-01-01", TS=4)
```

# Index

## \* datasets

CampinasRain, 2

lonlat, 3

Map, 3

rainTS4, 4

CampinasRain, 2

lonlat, 3

Map, 3

rainTS4, 4

SPIChanges, 5

TSaggreg, 6