

Package: Skillings.Mack (via r-universe)

August 20, 2024

Version 1.10

Date 2015-09-25

Title The Skillings-Mack Test Statistic for Block Designs with Missing Observations

Author Patchanok Srisuradetchai

Maintainer Patchanok Srisuradetchai <s.patchanok@gmail.com>

Depends R (>= 3.1.1)

Imports MASS, stats, matrixcalc

Description A generalization of the statistic used in Friedman's ANOVA method and in Durbin's rank test. This nonparametric statistical test is useful for the data obtained from block designs with missing observations occurring randomly. A resulting p-value is based on the chi-squared distribution and Monte Carlo method.

License GPL (>= 2)

LazyLoad yes

NeedsCompilation no

Repository CRAN

Date/Publication 2015-09-25 14:16:27

Contents

Ski.Mack	2
SkiMack	3

Index	6
--------------	----------

Ski.Mack

*The Skillings-Mack test statistic***Description**

The Skillings-Mack test statistic is a generalization of the statistic used in Friedman's ANOVA method and in Durbin's rank test. This nonparametric statistical test is useful for the data obtained from block designs with missing observations occurring randomly. The results are the Skillings-Mack statistic, p-values based on the chi-squared distribution and Monte Carlo method. The Monte Carlo method is recommended for approximating p-values when there are many ties and/or small designs with missing values are conducted.

Usage

```
Ski.Mack(y, groups=NULL, blocks=NULL, simulate.p.value = FALSE, B = 10000)
```

Arguments

y	Either a numeric vector of data values, or a data matrix. If a matrix is used, columns and rows are correspondent to blocks and treatments (groups), respectively.
groups	A vector containing group (treatment) indices for the corresponding y's which is a vector; this vector can be ignored if y is a matrix. Either a numeric or character vector is accepted.
blocks	A vector containing block indices for the corresponding y's; this vector can be ignored if y is a matrix. Either a numeric or character vector is accepted.
simulate.p.value	If TRUE, an estimated p-value based on the Monte Carlo method is calculated. The default is FALSE.
B	If simulate.p.value = TRUE, the default number of replications is 10,000.

Details

This function implements the Skilling-Mack test statistic for data obtained from block designs with missing observations. The `skillings.mack` function can also applied to balanced and partially balanced incomplete block designs. It reads a dataset in a format of a vector or a matrix and returns a p-value based on the chi-squared distribution. If `simulate.p.value = TRUE`, an estimated p-value based on Monte Carlo will be provided. If there are ties in a particular block, the average rank for those will be generated in simulating data under the null hypothesis.

Value

<code>\$nblocks</code>	a number of blocks in a design
<code>\$ntreatments</code>	a number of groups (treatments) in a design
<code>\$rawdata</code>	a matrix containing the data that was read from y

\$rankdata a matrix containing ranks of the data
 \$varCovarMatrix a variance-covariance matrix for adjusted treatment sums
 \$adjustedSum an adjusted treatment sum for the j th treatment

Author(s)

Patchanok Srisuradetchai <s.patchanok@gmail.com>

References

Skillings, J.H., Mack, G.A.: On the Use of a Friedman-Type Statistic in Balanced and Unbalanced Block Designs. *Technometrics* 23(2), 171-177 (1981)

Examples

```
## Skillings and Mack (1981), Table 1 page 173
## Comparison of four methods of assembling a product
B <- rep(c(1,2,3,4,5,6,7,8,9),rep(4,9))
G <- rep(c('A','B','C','D'),9)
y <- c(3.2,4.1,3.8,4.2, 3.1,3.9,3.4,4.0, 4.3,3.5,4.6,4.8,
      3.5,3.6,3.9,4.0, 3.6,4.2,3.7,3.9, 4.5,4.7,3.7, NA,
      NA ,4.2,3.4,NA , 4.3,4.6,4.4,4.9, 3.5, NA,3.7, 3.9)
Ski.Mack(y,groups = G,blocks = B)

## y is a matrix
maty <- matrix(
  c(3.2,4.1,3.8,4.2, 3.1,3.9,3.4,4.0, 4.3,3.5,4.6,4.8,
    3.5,3.6,3.9,4.0, 3.6,4.2,3.7,3.9, 4.5,4.7,3.7, NA,
    NA ,4.2,3.4,NA , 4.3,4.6,4.4,4.9, 3.5, NA,3.7, 3.9),
  ncol=9,byrow=FALSE)
Ski.Mack(maty, simulate.p.value = TRUE, B = 1000)
```

SkiMack

The Skillings-Mack test statistic

Description

The Skillings-Mack test statistic is a generalization of the statistic used in Friedman's ANOVA method and in Durbin's rank test. This nonparametric statistical test is useful for the data obtained from block designs with missing observations occurring randomly. The results from the SkiMack function are the Skillings-Mack statistic, p-values based on the chi-squared distribution and Monte Carlo method. The Monte Carlo method is recommended for approximating p-values when there are many ties and/or small designs with missing values are conducted.

Usage

```
SkiMack(y, groups=NULL, blocks=NULL, simulate.p.value = FALSE, B = 10000)
```

Arguments

y	Either a numeric vector of data values, or a data matrix. If a matrix is used, columns and rows are correspondent to blocks and treatments (groups), respectively.
groups	A vector giving the group (treatment) for the corresponding elements of y if this is a vector; ignored if y is a matrix. Either a numeric or character vector is accepted.
blocks	A vector giving the block for the corresponding elements of y if this is a vector; ignored if y is a matrix. Either a numeric or character vector is accepted.
simulate.p.value	If TRUE, an estimated p-value based on the Monte Carlo approach is calculated. The default is FALSE.
B	If simulate.p.value = TRUE, the default number of replications is 10,000.

Details

This function implements the Skilling-Mack test statistic for data obtained from block designs with missing observations. The `skillings.mack` function can also applied to balanced and partially balanced incomplete block designs. It reads a dataset from a vector or a matrix and returns a p-value based on the chi-squared distribution. If `simulate.p.value = TRUE`, an estimated p-value based on Monte Carlo will be provided. If there are ties in a particular block, the average rank for those will be generated in simulating data under the null hypothesis.

Value

<code>\$Nblocks</code>	the number of blocks in a design
<code>\$Ntreatments</code>	the number of groups (treatments) in a design
<code>\$rawdata</code>	a matrix containing the data that was read from y
<code>\$rankdata</code>	a matrix containing ranks of the data
<code>\$varCovarMatrix</code>	the variance-covariance matrix for adjusted treatment sums
<code>\$adjustedSum</code>	the adjusted treatment sum, A_j , for the j th treatment

Author(s)

Patchanok Srisuradetchai <s.patchanok@gmail.com>
 John J. Borkowski <jobo@math.montana.edu>

References

Skillings, J.H., Mack, G.A.: On the Use of a Friedman-Type Statistic in Balanced and Unbalanced Block Designs. *Technometrics* 23(2), 171-177 (1981)

Examples

```
## Skilling and Mack (1981), Table 1 page 173
## Comparison of four methods of assembling a product
B <- rep(c(1,2,3,4,5,6,7,8,9),rep(4,9))
G <- rep(c('A','B','C','D'),9)
y <- c(3.2,4.1,3.8,4.2, 3.1,3.9,3.4,4.0, 4.3,3.5,4.6,4.8,
      3.5,3.6,3.9,4.0, 3.6,4.2,3.7,3.9, 4.5,4.7,3.7, NA,
      NA ,4.2,3.4,NA , 4.3,4.6,4.4,4.9, 3.5, NA,3.7, 3.9)
SkiMack(y,groups = G,blocks = B)

## y is a matrix
maty <- matrix(
  c(3.2,4.1,3.8,4.2, 3.1,3.9,3.4,4.0, 4.3,3.5,4.6,4.8,
    3.5,3.6,3.9,4.0, 3.6,4.2,3.7,3.9, 4.5,4.7,3.7, NA,
    NA ,4.2,3.4,NA , 4.3,4.6,4.4,4.9, 3.5, NA,3.7, 3.9),
  ncol=9,byrow=FALSE)
SkiMack(maty, simulate.p.value = TRUE, B = 1000)
```

Index

* **Skilling-Mack, missing values**

SkiMack, [3](#)

* **block designs, missing values**

Ski.Mack, [2](#)

Ski.Mack, [2](#)

SkiMack, [3](#)