

Package: TDSTNN (via r-universe)

October 24, 2024

Type Package

Title Time Delay Spatio Temporal Neural Network

Version 0.1.0

Depends R (>= 4.2.3), nnet

Description STARMA (Space-Time Autoregressive Moving Average) models are commonly utilized in modeling and forecasting spatiotemporal time series data. However, the intricate nonlinear dynamics observed in many space-time rainfall patterns often exceed the capabilities of conventional STARMA models. This R package enables the fitting of Time Delay Spatio-Temporal Neural Networks, which are adept at handling such complex nonlinear dynamics efficiently. For detailed methodology, please refer to Saha et al. (2020) <[doi:10.1007/s00704-020-03374-2](https://doi.org/10.1007/s00704-020-03374-2)>.

Encoding UTF-8

License GPL-3

NeedsCompilation no

Author Mrinmoy Ray [aut, cre], Rajeev Ranjan Kumar [aut, ctb], Kanchan Sinha [aut, ctb], K. N. Singh [aut, ctb]

Maintainer Mrinmoy Ray <mrinmoy4848@gmail.com>

Repository CRAN

Date/Publication 2024-05-26 17:10:02 UTC

Contents

| | |
|----------------|----------|
| STNN | 2 |
| Index | 4 |

 STNN

Spatio-Temporal Neural Network

Description

The STNN function fits a Time-Delay Spatio-Temporal Neural Network model for space-time time series data.

Usage

```
STNN(data, lag, weight0, weight1, hs, h)
```

Arguments

| | |
|---------|--|
| data | Spatio-temporal time series (ts) data. |
| lag | Number of time lag(s). |
| weight0 | Zero-order weight matrix. |
| weight1 | First-order weight matrix. |
| hs | Number of hidden layer(s). |
| h | The forecast horizon. |

Details

This function enables you to apply the Time-delay Spatio-Temporal model for analyzing space-time series data.

Value

| | |
|-------------------|---|
| Model Summary | Summary of the fitted STNN |
| Fitted values | Fitted values of STNN |
| Forecasted values | h step ahead forecasted values employing STNN |

Author(s)

Mrinmoy Ray, Rajeev Ranjan Kumar, Kanchan Sinha, K. N. Singh

References

Saha, A., Singh, K.N., Ray, M. et al. A hybrid spatio-temporal modelling: an application to space-time rainfall forecasting. *Theor Appl Climatol* 142, 1271–1282 (2020).

See Also

ANN

Examples

```
ts.sim1 <- 50+arima.sim(list(order = c(1,1,0), ar = 0.7), n = 100)
ts.sim2<-70+arima.sim(list(order = c(1,1,0), ar = 0.8), n = 100)
weight0=diag(1, 2, 2)
weight1=matrix(c(0, 1, 1, 0), nrow= 2, ncol = 2, byrow = TRUE)
zz=as.matrix(cbind(ts.sim1,ts.sim2))
data=zz
lag=1
hs=2
h=5
STNN(data,lag,weight0,weight1,hs,h)
```

Index

* **STNN**
STNN, 2

STNN, 2