

Package: mable (via r-universe)

November 1, 2024

Title Maximum Approximate Bernstein/Beta Likelihood Estimation

Version 4.1.1

Date 2024-09-27

Author Zhong Guan [aut, cre]

Maintainer Zhong Guan <zguan@iu.edu>

Depends R (>= 3.5.0)

Description Fit data from a continuous population with a smooth density on finite interval by an approximate Bernstein polynomial model which is a mixture of certain beta distributions and find maximum approximate Bernstein likelihood estimator of the unknown coefficients. Consequently, maximum likelihood estimates of the unknown density, distribution functions, and more can be obtained. If the support of the density is not the unit interval then transformation can be applied. This is an implementation of the methods proposed by the author of this package published in the Journal of Nonparametric Statistics: Guan (2016) <doi:10.1080/10485252.2016.1163349> and Guan (2017) <doi:10.1080/10485252.2017.1374384>. For data with covariates, under some semiparametric regression models such as Cox proportional hazards model and the accelerated failure time model, the baseline survival function can be estimated smoothly based on general interval censored data.

License LGPL (>= 2.0, < 3)

LazyData true

Encoding UTF-8

Imports survival, graphics, stats, icenReg, parallel, doParallel, foreach, iterators, tcltk, quadprog, LowRankQP, mnormt, rlang

Suggests mixtools, Epi, ICsurv, interval, knitr, rmarkdown, pbapply, markdown, ks, multimode

BuildVignettes true

VignetteBuilder knitr

RoxygenNote 7.3.1

NeedsCompilation yes

Repository CRAN

Date/Publication 2024-10-01 08:40:02 UTC

Contents

chicken.embryo	3
chpt.exp	4
copula2d	4
copula2d.cond	5
corr.hellinger	6
cosmesis	7
dmixbeta	7
dmixmvbeta	9
dtmixbeta	10
mable	11
mable.aft	15
mable.copula	17
mable.ctrl	20
mable.decon	21
mable.dr	23
mable.dr.group	25
mable.group	27
mable.hellcorr	29
mable.ic	31
mable.mvar	33
mable.ph	36
mable.po	38
mable.reg	41
made.copula	43
made.density	44
made.mvar	45
madem.copula	47
madem.density	48
maple.aft	49
maple.dr	52
maple.dr.group	54
maple.ph	56
maple.po	58
marginal.p	61
mvecdf	62
mvpbeta	62
optim.gcp	63
optimable	64
pancreas	66
plot.mable	67

<i>chicken.embryo</i>	3
plot.mable_reg	68
se.coef.dr	69
summary.mable	70
umc.mat	70
Vaal.Flow	71
weib.gpo	71
Index	74

<i>chicken.embryo</i>	<i>Chicken Embryo Data</i>
-----------------------	----------------------------

Description

The chicken embryo dataset which contains day, number of days, and nT, the corresponding frequencies.

Usage

```
data(chicken.embryo)
```

Format

The format is: List of 2: day: int [1:21] 1 2 3 4 5 6 7 8 9 10 ...; nT : int [1:21] 6 5 11 2 2 3 0 0 0 0 ...

Source

Jassim, E. W., Grossman, M., Koops, W. J. And Luykx, R. A. J. (1996). Multi-phasic analysis of embryonic mortality in chickens. *Poultry Sci.* 75, 464-71.

References

Kuurman, W. W., Bailey, B. A., Koops, W. J. And Grossman, M. (2003). A model for failure of a chicken embryo to survive incubation. *Poultry Sci.* 82, 214-22.

Guan, Z. (2017) Bernstein polynomial model for grouped continuous data. *Journal of Nonparametric Statistics*, 29(4):831-848.

Examples

```
data(chicken.embryo)
```

chpt.exp

Exponential change-point

Description

Exponential change-point

Usage

```
chpt.exp(x)
```

Arguments

x a vector of nondecreasing values of log-likelihood or -log of distance

Value

a list of exponential change-point, its p-value, and the likelihood ratios of the exponential change-point model

copula2d

Some Bivariate Copulas

Description

Parametric bivariate copulas, densities, and random number generators

Usage

```
d2dcop.asym(u, v, lambda, copula = "clayton", ...)
```

```
p2dcop.asym(u, v, lambda, copula = "clayton", ...)
```

```
r2dcop.asym(n, lambda, copula = "clayton", ...)
```

```
dcopula(u, v, copula, ...)
```

```
pcopula(u, v, copula, ...)
```

```
rcopula(n, copula, ...)
```

Arguments

u, v	vectors of same length at which the copula and its density is evaluated
lambda	a vector of three mixing proportions which sum to one
copula	the name of a copula to be called or a base copula for constructing asymmetric copula(see Details)
...	the parameter(s) of copula, theta for most of the models, and df, the degrees of freedom if copula='t', or m if copula='nakagami'
n	number of random vectors to be generated

Details

The names of available copulas are 'amh' (Ali-Mikhail-Haq), 'bern' (Bernstein polynomial model), 'clayton' (Clayton), 'exponential' (Exponential), 'fgm' (Farlie-Gumbel-Morgenstern), 'frank' (Frank), 'gauss' (Gaussian), 'gumbel' (Gumbel), 'indep' (Independence), 'joe' (Joe), 'nakagami' (Nakagami-m), 'plackett' (Plackett), 't' (Student's t). `d2dcop.asym`, etc, calculate the constructive asymmetric copula of Wu (2014) using base copula C_θ with mixing proportions $p = (\lambda_1, \lambda_2, \lambda_3)$ and parameter values $\theta = (\theta_1, \theta_2, \theta_3)$: $\lambda_0 C_{\theta_0}(u, v) + \lambda_1 [v - C_{\theta_1}(1 - u, v)] + \lambda_2 [u - C_{\theta_2}(u, 1 - v)]$. If copula='t' or 'nakagami', df or m must be also given.

Value

a vector of copula or its density values evaluated at (u, v) or an n x 2 matrix of the generated observations

References

Nelsen, R. B. (1999). *An Introduction to Copulas*. Springer Series in Statistics. New York: Springer. Wu, S. (2014). Construction of asymmetric copulas and its application in two-dimensional reliability modelling. *European Journal of Operational Research* 238 (2), 476–485.

 copula2d.cond

Some Parametric Conditional Bivariate Copulas

Description

Density, distribution function, quantile function and random generation for conditional copula $C(u|V = v)$ of U given $V = v$ related to parametric bivariate copula $C(u, v) = P(U \leq u, V \leq v)$.

Usage

```
dcopula.cond(u, v, copula, ...)
```

```
pcopula.cond(u, v, copula, ...)
```

```
qcopula.cond(p, v, copula, ...)
```

```
rcopula.cond(n, v, copula, ...)
```

Arguments

u	vector of U values at which the copula density is evaluated
v	a given value of V under which the conditional copula and its density is evaluated
copula	the name of a copula density to be called (see Details)
...	the parameter(s) of copula
p	a vector of probabilities
n	number of observations to be generated from conditional copula $C(u V = v)$.

Details

the names of available copulas are 'amh' (Ali-Mikhail-Haq), 'bern' (Bernstein polynomial model), 'clayton' (Clayton), 'exponential' (Exponential), 'fgm' (Farlie-Gumbel-Morgenstern), 'frank' (Frank), 'gauss' (Gaussian), 'gumbel' (Gumbel), 'indep' (Independence), 'joe' (Joe), 'nakagami' (Nakagami-m), 'plackett' (Plackett), 't' (Student's t).

Value

a vector of copula density values evaluated at u given $V=v$ or a vector of n generated u values from conditional copula $C(u|V = v)$.

corr.hellinger	<i>Bhattacharyya coefficient and Hellinger correlation</i>
----------------	--

Description

Bhattacharyya coefficient and Hellinger correlation

Usage

```
corr.hellinger(dcopula, ...)
```

Arguments

dcopula	a function object defining a 2d copula density function
...	argument(s) of copula density function

Value

Bhattacharyya coefficient B and Hellinger correlation eta

References

Geenens, G. and Lafaye de Micheaux, P. (2022). The Hellinger correlation. *Journal of the American Statistical Association* 117(538), 639–653.

`cosmesis`*Breast cosmesis data*

Description

Data contain the interval-censored times to cosmetic deterioration for breast cancer patients undergoing radiation or radiation plus chemotherapy.

Usage

```
data(cosmesis)
```

Format

A data frame with 94 observations on the following 3 variables.

- left left endpoint of the censoring interval in months
- right right endpoint of the censoring interval in months
- treat a factor with levels RT and RCT representing radiotherapy-only and radiation plus chemotherapy treatments, respectively

Source

Finkelstein, D. M. and Wolfe, R. A. (1985) A semiparametric model for regression analysis of interval-censored failure time data. *Biometrics* 41, 933–945.

References

Finkelstein, D. M. (1986) A proportional hazards model for interval-censored failure time data. *Biometrics* 42, 845–854.

Examples

```
data(cosmesis)
```

`dmixbeta`*Mixture Beta Distribution*

Description

Density, distribution function, quantile function and pseudorandom number generation for the Bernstein polynomial model, mixture of beta distributions, with shapes $(i + 1, m - i + 1)$, $i = 0, \dots, m$, given mixture proportions $p = (p_0, \dots, p_m)$ and support interval.

Usage

```
dmixbeta(x, p, interval = c(0, 1))
```

```
pmixbeta(x, p, interval = c(0, 1))
```

```
qmixbeta(u, p, interval = c(0, 1))
```

```
rmixbeta(n, p, interval = c(0, 1))
```

Arguments

x	a vector of quantiles
p	a vector of m+1 values. The m+1 components of p must be nonnegative and sum to one for mixture beta distribution. See 'Details'.
interval	support/truncation interval [a, b].
u	a vector of probabilities
n	sample size

Details

The density of the mixture beta distribution on an interval $[a, b]$ can be written as a Bernstein polynomial $f_m(x; p) = (b - a)^{-1} \sum_{i=0}^m p_i \beta_{mi}[(x - a)/(b - a)]/(b - a)$, where $p = (p_0, \dots, p_m)$, $p_i \geq 0$, $\sum_{i=0}^m p_i = 1$ and $\beta_{mi}(u) = (m + 1) \binom{m}{i} u^i (1 - u)^{m-i}$, $i = 0, 1, \dots, m$, is the beta density with shapes $(i + 1, m - i + 1)$. The cumulative distribution function is $F_m(x; p) = \sum_{i=0}^m p_i B_{mi}[(x - a)/(b - a)]$, where $B_{mi}(u)$, $i = 0, 1, \dots, m$, is the beta cumulative distribution function with shapes $(i + 1, m - i + 1)$. If $\pi = \sum_{i=0}^m p_i < 1$, then f_m/π is a truncated density on $[a, b]$ with cumulative distribution function F_m/π . The argument p may be any numeric vector of m+1 values when pmixbeta() and qmixbeta() return the integral function $F_m(x; p)$ and its inverse, respectively, and dmixbeta() returns a Bernstein polynomial $f_m(x; p)$. If components of p are not all nonnegative or do not sum to one, warning message will be returned.

Value

A vector of $f_m(x; p)$ or $F_m(x; p)$ values at x . dmixbeta returns the density, pmixbeta returns the cumulative distribution function, qmixbeta returns the quantile function, and rmixbeta generates pseudo random numbers.

Author(s)

Zhong Guan <zguan@iu.edu>

References

- Bernstein, S.N. (1912), Demonstration du theoreme de Weierstrass fondee sur le calcul des probabilities, Communications of the Kharkov Mathematical Society, 13, 1–2.
- Guan, Z. (2016) Efficient and robust density estimation using Bernstein type polynomials. *Journal of Nonparametric Statistics*, 28(2):250-271.

Guan, Z. (2017) Bernstein polynomial model for grouped continuous data. *Journal of Nonparametric Statistics*, 29(4):831-848.

See Also

[mable](#)

Examples

```
# classical Bernstein polynomial approximation
a<-4; b<-4; m<-200
x<-seq(a,b,len=512)
u<-(0:m)/m
p<-dnorm(a+(b-a)*u)
plot(x, dnorm(x), type="l")
lines(x, (b-a)*dmixbeta(x, p, c(a, b))/(m+1), lty=2, col=2)
legend(a, dnorm(0), lty=1:2, col=1:2, c(expression(f(x)==phi(x)),
expression(B^{f}(x))))
```

dmixmvbeta

Multivariate Mixture Beta Distribution

Description

Density, distribution function, and pseudorandom number generation for the multivariate Bernstein polynomial model, mixture of multivariate beta distributions, with given mixture proportions $p = (p_0, \dots, p_{K-1})$, given degrees $m = (m_1, \dots, m_d)$, and support interval.

Usage

```
dmixmvbeta(x, p, m, interval = NULL)
```

```
pmixmvbeta(x, p, m, interval = NULL)
```

```
rmixmvbeta(n, p, m, interval = NULL)
```

Arguments

x	a matrix with d columns or a vector of length d within support hyperrectangle $[a, b] = [a_1, b_1] \times \dots \times [a_d, b_d]$
p	a vector of K values. All components of p must be nonnegative and sum to one for the mixture multivariate beta distribution. See 'Details'.
m	a vector of degrees, (m_1, \dots, m_d)
interval	a vector of two endpoints or a 2 x d matrix, each column containing the endpoints of support/truncation interval for each marginal density. If missing, the i-th column is assigned as $c(\theta, 1)$.
n	sample size

Details

dmixmvbeta() returns a linear combination f_m of d -variate beta densities on $[a, b]$, $\beta_{mj}(x) = \prod_{i=1}^d \beta_{m_i, j_i}[(x_i - a_i)/(b_i - a_i)]/(b_i - a_i)$, with coefficients $p(j_1, \dots, j_d)$, $0 \leq j_i \leq m_i$, $i = 1, \dots, d$, where $[a, b] = [a_1, b_1] \times \dots \times [a_d, b_d]$ is a hyperrectangle, and the coefficients are arranged in the column-major order of $j = (j_1, \dots, j_d)$, p_0, \dots, p_{K-1} , where $K = \prod_{i=1}^d (m_i + 1)$. pmixmvbeta() returns a linear combination F_m of the distribution functions of d -variate beta distribution.

If all p_i 's are nonnegative and sum to one, then p are the mixture proportions of the mixture multivariate beta distribution.

 dtmixbeta

Exponentially Tilted Mixture Beta Distribution

Description

Density, distribution function, quantile function and pseudorandom number generation for the exponentially tilted mixture of beta distributions, with shapes $(i + 1, m - i + 1)$, $i = 0, \dots, m$, given mixture proportions $p = (p_0, \dots, p_m)$ and support interval.

Usage

```
dtmixbeta(x, p, alpha, interval = c(0, 1), regr, ...)
```

```
ptmixbeta(x, p, alpha, interval = c(0, 1), regr, ...)
```

```
qtmixbeta(u, p, alpha, interval = c(0, 1), regr, ...)
```

```
rtmixbeta(n, p, alpha, interval = c(0, 1), regr, ...)
```

Arguments

x	a vector of quantiles
p	a vector of $m+1$ components of p must be nonnegative and sum to one for mixture beta distribution. See 'Details'.
alpha	regression coefficients
interval	support/truncation interval $[a, b]$.
regr	regressor vector function $r(x) = (1, r_1(x), \dots, r_d(x))$ which returns $n \times (d+1)$ matrix, $n = \text{length}(x)$
...	additional arguments to be passed to regr
u	a vector of probabilities
n	sample size

Details

The density of the mixture exponentially tilted beta distribution on an interval $[a, b]$ can be written $f_m(x; p) = (b-a)^{-1} \exp(\alpha' r(x)) \sum_{i=0}^m p_i \beta_{mi}[(x-a)/(b-a)]/(b-a)$, where $p = (p_0, \dots, p_m)$, $p_i \geq 0$, $\sum_{i=0}^m p_i = 1$ and $\beta_{mi}(u) = (m+1) \binom{m}{i} u^i (1-u)^{m-i}$, $i = 0, 1, \dots, m$, is the beta density with shapes $(i+1, m-i+1)$. The cumulative distribution function is $F_m(x; p) = \sum_{i=0}^m p_i B_{mi}[(x-a)/(b-a); \alpha]$, where $B_{mi}(u; \alpha)$, $i = 0, 1, \dots, m$, is the exponentially tilted beta cumulative distribution function with shapes $(i+1, m-i+1)$.

Value

A vector of $f_m(x; p)$ or $F_m(x; p)$ values at x . `dmixbeta` returns the density, `pmixbeta` returns the cumulative distribution function, `qmixbeta` returns the quantile function, and `rmixbeta` generates pseudo random numbers.

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z., Application of Bernstein Polynomial Model to Density and ROC Estimation in a Semi-parametric Density Ratio Model

See Also

[mable](#)

Examples

```
# classical Bernstein polynomial approximation
a<-4; b<-4; m<-200
x<-seq(a,b,len=512)
u<-(0:m)/m
p<-dnorm(a+(b-a)*u)
plot(x, dnorm(x), type="l")
lines(x, (b-a)*dmixbeta(x, p, c(a, b))/(m+1), lty=2, col=2)
legend(a, dnorm(0), lty=1:2, col=1:2, c(expression(f(x)==phi(x)),
expression(B^{f}(x))))
```

mable

Maximum Approximate Bernstein Likelihood Estimate of Univariate or Multivariate Density Function

Description

Maximum approximate Bernstein/Beta likelihood estimation based on one-sample raw data with an optimal selected by the change-point method among $m_0:m_1$ or a preselected model degree m .

Usage

```
mable(
  x,
  M,
  interval = 0:1,
  IC = c("none", "aic", "hqic", "all"),
  vb = 0,
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

x	a (non-empty) numeric n-vector or n x d matrix or data.frame of n observations.
M	a positive integer or a vector (m0, m1). If M = m or m0 = m1 = m, then m is a preselected degree. If m0 < m1 it specifies the set of consecutive candidate model degrees m0:m1 for searching an optimal degree, where m1 - m0 > 3.
interval	a vector containing the endpoints of supporting/truncation interval c(a, b)
IC	information criterion(s) in addition to Bayesian information criterion (BIC). Current choices are "aic" (Akaike information criterion) and/or "hqic" (Hannan-Quinn information criterion).
vb	code for vanishing boundary constraints, -1: f0(a)=0 only, 1: f0(b)=0 only, 2: both, 0: none (default).
controls	Object of class mable.ctrl() specifying iteration limit and the convergence criterion eps. Default is mable.ctrl. See Details.
progress	if TRUE a text progressbar is displayed

Details

Any continuous density function f on a known closed supporting interval $[a, b]$ can be estimated by Bernstein polynomial $f_m(x; p) = \sum_{i=0}^m p_i \beta_{mi}[(x-a)/(b-a)]/(b-a)$, where $p = (p_0, \dots, p_m)$, $p_i \geq 0$, $\sum_{i=0}^m p_i = 1$ and $\beta_{mi}(u) = (m+1) \binom{m}{i} u^i (1-u)^{m-i}$, $i = 0, 1, \dots, m$, is the beta density with shapes $(i+1, m-i+1)$. For each m , the MABLE of the coefficients p , the mixture proportions, are obtained using EM algorithm. The EM iteration for each candidate m stops if either the total absolute change of the log likelihood and the coefficients of Bernstein polynomial is smaller than `eps` or the maximum number of iterations `maxit` is reached.

If $m_0 < m_1$, an optimal model degree is selected as the change-point of the increments of log-likelihood, log likelihood ratios, for $m \in \{m_0, m_0 + 1, \dots, m_1\}$. Alternatively, one can choose an optimal degree based on the BIC (Schwarz, 1978) which are evaluated at $m \in \{m_0, m_0 + 1, \dots, m_1\}$. The search for optimal degree m is stopped if either m_1 is reached with a warning or the test for change-point results in a p-value `pval` smaller than `sig.level`. The BIC for a given degree m is calculated as in Schwarz (1978) where the dimension of the model is $d = \#\{i : \hat{p}_i \geq \epsilon, i = 0, \dots, m\} - 1$ and a default ϵ is chosen as `.Machine$double.eps`.

If data show a clearly multimodal distribution by plotting the histogram for example, the model degree is usually large. The range `M` should be large enough to cover the optimal degree and the

computation is time-consuming. In this case the iterative method of moment with an initial selected by a method of mode which is implemented by `optimable` can be used to reduce the computation time.

Value

A list with components

- `m` the given or a selected degree by method of change-point
- `p` the estimated vector of mixture proportions $p = (p_0, \dots, p_m)$ with the selected/given optimal degree `m`
- `mloglik` the maximum log-likelihood at degree `m`
- `interval` support/truncation interval (a, b)
- `convergence` An integer code. 0 indicates successful completion (all the EM iterations are convergent and an optimal degree is successfully selected in `M`). Possible error codes are
 - 1, indicates that the iteration limit `maxit` had been reached in at least one EM iteration;
 - 2, the search did not finish before `m1`.
- `delta` the convergence criterion `delta` value

and, if `m0 < m1`,

- `M` the vector (m_0, m_1) , where `m1`, if greater than `m0`, is the largest candidate when the search stopped
- `lk` log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- `lr` likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- `ic` a list containing the selected information criterion(s)
- `pval` the p-values of the change-point tests for choosing optimal model degree
- `chpts` the change-points chosen with the given candidate model degrees

Note

Since the Bernstein polynomial model of degree m is nested in the model of degree $m + 1$, the maximum likelihood is increasing in m . The change-point method is used to choose an optimal degree m . The degree can also be chosen by a method of moment and a method of mode which are implemented by function `optimal()`.

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. (2016) Efficient and robust density estimation using Bernstein type polynomials. *Journal of Nonparametric Statistics*, 28(2):250-271. Wang, T. and Guan, Z.,(2019) Bernstein Polynomial Model for Nonparametric Multivariate Density, *Statistics*, Vol. 53, no. 2, 321-338

See Also[optimable](#)**Examples**

```

# Vaal Rive Flow Data
data(Vaal.Flow)
x<-Vaal.Flow$Flow
res<-mable(x, M = c(2,100), interval = c(0, 3000), controls =
  mable.ctrl(sig.level = 1e-8, maxit = 2000, eps = 1.0e-9))
op<-par(mfrow = c(1,2),lwd = 2)
layout(rbind(c(1, 2), c(3, 3)))
plot(res, which = "likelihood", cex = .5)
plot(res, which = c("change-point"), lgd.x = "topright")
hist(x, prob = TRUE, xlim = c(0,3000), ylim = c(0,.0022), breaks = 100*(0:30),
  main = "Histogram and Densities of the Annual Flow of Vaal River",
  border = "dark grey",lwd = 1,xlab = "x", ylab = "f(x)", col = "light grey")
lines(density(x, bw = "nrd0", adjust = 1), lty = 4, col = 4)
lines(y<-seq(0, 3000, length = 100), dlnorm(y, mean(log(x)),
  sqrt(var(log(x))))), lty = 2, col = 2)
plot(res, which = "density", add = TRUE)
legend("top", lty = c(1, 2, 4), col = c(1, 2, 4), bty = "n",
c(expression(paste("MABLE: ",hat(f)[B])),
  expression(paste("Log-Normal: ",hat(f)[P])),
  expression(paste("KDE: ",hat(f)[K]))))
par(op)

# Old Faithful Data
library(mixtools)
x<-faithful$eruptions
a<-0; b<-7
v<-seq(a, b,len = 512)
mu<-c(2,4.5); sig<-c(1,1)
pmix<-normalmixEM(x,.5, mu, sig)
lam<-pmix$lambda; mu<-pmix$mu; sig<-pmix$sigma
y1<-lam[1]*dnorm(v,mu[1], sig[1])+lam[2]*dnorm(v, mu[2], sig[2])
res<-mable(x, M = c(2,300), interval = c(a,b), controls =
  mable.ctrl(sig.level = 1e-8, maxit = 2000L, eps = 1.0e-7))
op<-par(mfrow = c(1,2),lwd = 2)
layout(rbind(c(1, 2), c(3, 3)))
plot(res, which = "likelihood")
plot(res, which = "change-point")
hist(x, breaks = seq(0,7.5,len = 20), xlim = c(0,7), ylim = c(0,.7),
  prob = TRUE,xlab = "t", ylab = "f(t)", col = "light grey",
  main = "Histogram and Density of
  Duration of Eruptions of Old Faithful")
lines(density(x, bw = "nrd0", adjust = 1), lty = 4, col = 4, lwd = 2)
plot(res, which = "density", add = TRUE)
lines(v, y1, lty = 2, col = 2, lwd = 2)
legend("topright", lty = c(1,2,4), col = c(1,2,4), lwd = 2, bty = "n",
  c(expression(paste("MABLE: ",hat(f)[B](x))),

```

```

expression(paste("Mixture: ",hat(f)[P](t))),
expression(paste("KDE: ",hat(f)[K](t))))
par(op)

```

mable.aft

Mable fit of Accelerated Failure Time Model

Description

Maximum approximate Bernstein/Beta likelihood estimation for accelerated failure time model based on interval censored data.

Usage

```

mable.aft(
  formula,
  data,
  M,
  g = NULL,
  p = NULL,
  tau = NULL,
  x0 = NULL,
  controls = mable.ctrl(),
  progress = TRUE
)

```

Arguments

formula	regression formula. Response must be cbind. See 'Details'.
data	a data frame containing variables in formula.
M	a positive integer or a vector (m_0, m_1). If $M = m_0$ or $m_0 = m_1 = m$, then m_0 is a preselected degree. If $m_0 < m_1$ it specifies the set of consecutive candidate model degrees $m_0:m_1$ for searching an optimal degree, where $m_1 - m_0 > 3$.
g	a d -vector of regression coefficients, default is the zero vector.
p	an initial coefficients of Bernstein polynomial of degree m_0 , default is the uniform initial.
tau	the right endpoint of the support or truncation interval $[0, \tau)$ of the baseline density. Default is NULL (unknown), otherwise if tau is given then it is taken as a known value of τ . See 'Details'.
x0	a data frame specifying working baseline covariates on the right-hand-side of formula. See 'Details'.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

Consider the accelerated failure time model with covariate for interval-censored failure time data: $S(t|x) = S(t \exp(\gamma^T(x - x_0))|x_0)$, where x and x_0 may contain dummy variables and interaction terms. The working baseline x_0 in arguments contains only the values of terms excluding dummy variables and interaction terms in the right-hand-side of formula. Thus g is the initial guess of the coefficients γ of $x - x_0$ and could be longer than x_0 . Let $f(t|x)$ and $F(t|x) = 1 - S(t|x)$ be the density and cumulative distribution functions of the event time given $X = x$, respectively. Then $f(t|x_0)$ on a truncation interval $[0, \tau]$ can be approximated by $f_m(t|x_0; p) = \tau^{-1} \sum_{i=0}^m p_i \beta_{mi}(t/\tau)$, where $p_i \geq 0$, $i = 0, \dots, m$, $\sum_{i=0}^m p_i = 1$, $\beta_{mi}(u)$ is the beta density with shapes $i + 1$ and $m - i + 1$, and τ is larger than the largest observed time, either uncensored time, or right endpoint of interval/left censored, or left endpoint of right censored time. So we can approximate $S(t|x_0)$ on $[0, \tau]$ by $S_m(t|x_0; p) = \sum_{i=0}^m p_i \bar{B}_{mi}(t/\tau)$, where $\bar{B}_{mi}(u)$ is the beta survival function with shapes $i + 1$ and $m - i + 1$.

Response variable should be of the form `cbind(1, u)`, where $(1, u)$ is the interval containing the event time. Data is uncensored if $1 = u$, right censored if $u = \text{Inf}$ or $u = \text{NA}$, and left censored data if $1 = 0$. The truncation time `tau` and the baseline `x0` should be chosen so that $S(t|x) = S(t \exp(\gamma^T(x - x_0))|x_0)$ on $[\tau, \infty)$ is negligible for all the observed x .

The search for optimal degree `m` stops if either `m1` is reached or the test for change-point results in a `p-value` `pval` smaller than `sig.level`.

Value

A list with components

- `m` the given or selected optimal degree `m`
- `p` the estimate of $p = (p_0, \dots, p_m)$, the coefficients of Bernstein polynomial of degree `m`
- `coefficients` the estimated regression coefficients of the AFT model
- `SE` the standard errors of the estimated regression coefficients
- `z` the z-scores of the estimated regression coefficients
- `mloglik` the maximum log-likelihood at an optimal degree `m`
- `tau.n` maximum observed time τ_n
- `tau` right endpoint of truncation interval $[0, \tau)$
- `x0` the working baseline covariates
- `egx0` the value of $e^{\gamma^T x_0}$
- `convergence` an integer code: 0 indicates a successful completion; 1 indicates that the search of an optimal degree using change-point method reached the maximum candidate degree; 2 indicates that the maximum iterations was reached for calculating \hat{p} and $\hat{\gamma}$ with the selected degree `m`, or the divergence of the last EM-like iteration for p or the divergence of the last (quasi) Newton iteration for γ ; 3 indicates 1 and 2.
- `delta` the final `delta` if `m0 = m1` or the final `pval` of the change-point for searching the optimal degree `m`;

and, if `m0 < m1`,

- `M` the vector (m_0, m_1) , where `m1` is the last candidate when the search stopped

- lk log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- lr likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- pval the p-values of the change-point tests for choosing optimal model degree
- chpts the change-points chosen with the given candidate model degrees

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. (2019) Maximum Approximate Likelihood Estimation in Accelerated Failure Time Model for Interval-Censored Data, arXiv:1911.07087.

See Also

[maple.aft](#)

Examples

```
## Breast Cosmesis Data
g <- 0.41 #Hanson and Johnson 2004, JCGS
aft.res<-mable.aft(cbind(left, right)~treat, data=cosmesis, M=c(1, 30),
                  g=g, tau=100, x0=data.frame(treat="RCT"))
op<-par(mfrow=c(1,2), lwd=1.5)
plot(x=aft.res, which="likelihood")
plot(x=aft.res, y=data.frame(treat="RT"), which="survival", model='aft', type="l", col=1,
     add=FALSE, main="Survival Function")
plot(x=aft.res, y=data.frame(treat="RCT"), which="survival", model='aft', lty=2, col=1)
legend("bottomleft", bty="n", lty=1:2, col=1, c("Radiation Only", "Radiation and Chemotherapy"))
par(op)
```

mable.copula

Maximum Approximate Bernstein Likelihood Estimate of Copula Density Function

Description

Maximum Approximate Bernstein Likelihood Estimate of Copula Density Function

Usage

```
mable.copula(
  x,
  M0 = 1,
  M,
  unif.mar = TRUE,
```

```

pseudo.obs = c("empirical", "mable"),
interval = NULL,
search = TRUE,
mar.deg = FALSE,
high.dim = FALSE,
controls = mable.ctrl(sig.level = 0.05),
progress = TRUE
)

```

Arguments

<code>x</code>	an $n \times d$ matrix or data.frame of multivariate sample of size n from d -variate distribution with hyperrectangular specified by <code>interval</code> .
<code>M0</code>	a nonnegative integer or a vector of d nonnegative integers specify starting candidate degrees for searching optimal degrees.
<code>M</code>	a positive integer or a vector of d positive integers specify the maximum candidate or the given model degrees for the joint density.
<code>unif.mar</code>	logical, whether all the marginals distributions are uniform or not. If not the pseudo observations will be created using empirical or mable marginal distributions.
<code>pseudo.obs</code>	"empirical": use empirical distribution to create pseudo, observations, or "mable": use mable of marginal cdfs to create pseudo observations
<code>interval</code>	a vector of two endpoints or a $2 \times d$ matrix, each column containing the endpoints of support/truncation interval for each marginal density. If missing, the i -th column is assigned as <code>extendrange(x[, i])</code> . If <code>unif.mar=TRUE</code> , then it is $[0, 1]^d$.
<code>search</code>	logical, whether to search optimal degrees between <code>M0</code> and <code>M</code> or not but use <code>M</code> as the given model degrees for the joint density.
<code>mar.deg</code>	logical, if TRUE (default), the optimal degrees are selected based on marginal data, otherwise, the optimal degrees are chosen by the method of change-point. See details.
<code>high.dim</code>	logical, data are high dimensional/large sample or not if TRUE, run a slower version procedure which requires less memory
<code>controls</code>	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion eps. Default is <code>mable.ctrl</code> . See Details.
<code>progress</code>	if TRUE a text progressbar is displayed

Details

A d -variate copula density $c(u)$ on $[0, 1]^d$ can be approximated by a mixture of d -variate beta densities on $[0, 1]^d$, $\beta_{mj}(x) = \prod_{i=1}^d \beta_{m_i, j_i}(u_i)$, with proportion $p(j_1, \dots, j_d)$, $0 \leq j_i \leq m_i, i = 1, \dots, d$, which satisfy the uniform marginal constraints, the copula (density) has uniform marginal cdf (pdf). If `search=TRUE` and `mar.deg=TRUE`, then the optimal degrees are $(\tilde{m}_1, \dots, \tilde{m}_d)$, where \tilde{m}_i is chosen based on marginal data of u_i , $i = 1, \dots, d$. If `search=TRUE` and `mar.deg=FALSE`, then the optimal degrees $(\hat{m}_1, \dots, \hat{m}_d)$ are chosen using a change-point method based on the joint data.

For large data and high dimensional density, the search for the model degrees might be time-consuming. Thus patience is needed.

Value

A list with components

- `m` a vector of the selected optimal degrees by the method of change-point
- `p` a vector of the mixture proportions $p(j_1, \dots, j_d)$, arranged in the column-major order of $j = (j_1, \dots, j_d)$, $0 \leq j_i \leq m_i, i = 1, \dots, d$.
- `mloglik` the maximum log-likelihood at an optimal degree `m`
- `pval` the p-values of change-points for choosing the optimal degrees for the marginal densities
- `M` the vector (`m1`, `m2`, . . . , `md`) at which the search of model degrees stopped. If `mar.deg=TRUE` `mi` is the largest candidate degree when the search stopped for the *i*-th marginal density
- `convergence` An integer code. 0 indicates successful completion(the EM iteration is convergent). 1 indicates that the iteration limit `maxit` had been reached in the EM iteration;
- if `unif.mar=FALSE`, `margin` contains objects of the results of `mable` fit to the marginal data

Author(s)

Zhong Guan <zguan@iu.edu>

References

Wang, T. and Guan, Z. (2019). Bernstein polynomial model for nonparametric multivariate density. *Statistics* 53(2), 321–338. Guan, Z., Nonparametric Maximum Likelihood Estimation of Copula

See Also

[mable](#), [mable.mvar](#)

Examples

```
## Simulated bivariate data from Gaussian copula

set.seed(1)
rho<-0.4; n<-1000
x<-rnorm(n)
u<-pnorm(cbind(rnorm(n, mean=rho*x, sd=sqrt(1-rho^2)),x))
res<- mable.copula(u, M = c(3,3), search =FALSE, mar.deg=FALSE, progress=FALSE)
plot(res, which="density")
```

`mable.ctrl`*Control parameters for mable fit*

Description

Control parameters for mable fit

Usage

```
mable.ctrl(  
  sig.level = 0.01,  
  eps = 1e-07,  
  maxit = 5000L,  
  eps.em = 1e-07,  
  maxit.em = 5000L,  
  eps.nt = 1e-07,  
  maxit.nt = 100L,  
  tini = 1e-04  
)
```

Arguments

<code>sig.level</code>	the significance level for change-point method of choosing optimal model degree
<code>eps</code>	convergence criterion for iteration involves EM like and Newton-Raphson iterations
<code>maxit</code>	maximum number of iterations involve EM like and Newton-Raphson iterations
<code>eps.em</code>	convergence criterion for EM like iteration
<code>maxit.em</code>	maximum number of EM like iterations
<code>eps.nt</code>	convergence criterion for Newton-Raphson iteration
<code>maxit.nt</code>	maximum number of Newton-Raphson iterations
<code>tini</code>	a small positive number used to make sure initial p is in the interior of the simplex

Value

a list of the arguments' values

Author(s)

Zhong Guan <zguan@iu.edu>

mable.decon

Mable deconvolution with a known error density

Description

Maximum approximate Bernstein/Beta likelihood estimation in additive density deconvolution model with a known error density.

Usage

```
mable.decon(
  y,
  gn = NULL,
  ...,
  M,
  interval = c(0, 1),
  IC = c("none", "aic", "hqic", "all"),
  vanished = TRUE,
  controls = mable.ctrl(maxit.em = 1e+05, eps.em = 1e-05, maxit.nt = 100, eps.nt = 1e-10),
  progress = TRUE
)
```

Arguments

y	vector of observed data values
gn	error density function if known, default is NULL if unknown
...	additional arguments to be passed to gn
M	a vector (m0, m1) specifies the set of consecutive candidate model degrees, M = m0:m1. If gn is unknown then M a 2 x 2 matrix whose rows (m0, m1) and (k0, k1) specify lower and upper bounds for degrees m and k, respectively.
interval	a finite vector (a, b), the endpoints of supporting/truncation interval if gn is known. Otherwise, it is a 2 x 2 matrix whose rows (a, b) and (a1, b1) specify supporting/truncation intervals of X and ε, respectively. See Details.
IC	information criterion(s) in addition to Bayesian information criterion (BIC). Current choices are "aic" (Akaike information criterion) and/or "hqic" (Hannan–Quinn information criterion).
vanished	logical whether the unknown error density vanishes at both end-points of [a1, b1]
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

Consider the additive measurement error model $Y = X + \epsilon$, where X has an unknown distribution F on a known support $[a, b]$, ϵ has a known or unknown distribution G , and X and ϵ are independent. We want to estimate density $f = F'$ based on independent observations, $y_i = x_i + \epsilon_i$, $i = 1, \dots, n$, of Y . We approximate f by a Bernstein polynomial model on $[a, b]$. If $g = G'$ is unknown on a known support $[a_1, b_1]$, then we approximate g by a Bernstein polynomial model on $[a_1, b_1]$, $a_1 < 0 < b_1$. We assume $E(\epsilon) = 0$. AIC and BIC methods are used to select model degrees (m, k) .

Value

A mable class object with components, if g is known,

- M the vector (m_0, m_1) , where m_1 is the last candidate degree when the search stopped
- m the selected optimal degree m
- p the estimate of $p = (p_0, \dots, p_m)$, the coefficients of Bernstein polynomial of degree m
- lk log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- lr likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- $convergence$ An integer code. 0 indicates an optimal degree is successfully selected in M . 1 indicates that the search stopped at m_1 .
- ic a list containing the selected information criterion(s)
- $pval$ the p-values of the change-point tests for choosing optimal model degree
- $chpts$ the change-points chosen with the given candidate model degrees

if g is unknown,

- M the 2 x 2 matrix with rows (m_0, m_1) and (k_0, k_1)
- nu_aic the selected optimal degrees (m, k) using AIC method
- p_aic the estimate of $p = (p_0, \dots, p_m)$, the coefficients of Bernstein polynomial model for f of degree m as in nu_aic
- q_aic the estimate of $q = (q_0, \dots, q_k)$, the coefficients of Bernstein polynomial model for g of degree k as in nu_aic
- nu_bic the selected optimal degrees (m, k) using BIC method
- p_bic the estimate of $p = (p_0, \dots, p_m)$, the coefficients of Bernstein polynomial model for f of degree m as in nu_bic
- q_bic the estimate of $q = (q_0, \dots, q_k)$, the coefficients of Bernstein polynomial model for g of degree k as in nu_bic
- lk matrix of log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$ and $k \in \{k_0, \dots, k_1\}$
- aic a matrix containing the Akaike information criterion(s) at $m \in \{m_0, \dots, m_1\}$ and $k \in \{k_0, \dots, k_1\}$
- bic a matrix containing the Bayesian information criterion(s) at $m \in \{m_0, \dots, m_1\}$ and $k \in \{k_0, \dots, k_1\}$

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z., (2019) Fast Nonparametric Maximum Likelihood Density Deconvolution Using Bernstein Polynomials, *Statistica Sinica*, doi:10.5705/ss.202018.0173

Examples

```
# A simulated normal dataset
set.seed(123)
mu<-1; sig<-2; a<-mu-sig*5; b<-mu+sig*5;
gn<-function(x) dnorm(x, 0, 1)
n<-50;
x<-rnorm(n, mu, sig); e<-rnorm(n); y<-x+e;
res<-mable.decon(y, gn, interval = c(a, b), M = c(5, 50))
op<-par(mfrow = c(2, 2),lwd = 2)
plot(res, which="likelihood")
plot(res, which="change-point", lgd.x="topright")
plot(xx<-seq(a, b, length=100), yy<-dnorm(xx, mu, sig), type="l", xlab="x",
      ylab="Density", ylim=c(0, max(yy)*1.1))
plot(res, which="density", types=c(2,3), colors=c(2,3))
# kernel density based on pure data
lines(density(x), lty=4, col=4)
legend("topright", bty="n", lty=1:4, col=1:4,
c(expression(f), expression(hat(f)[cp]), expression(hat(f)[bic]), expression(tilde(f)[K])))
plot(xx, yy<-pnorm(xx, mu, sig), type="l", xlab="x", ylab="Distribution Function")
plot(res, which="cumulative", types=c(2,3), colors=c(2,3))
legend("bottomright", bty="n", lty=1:3, col=1:3,
      c(expression(F), expression(hat(F)[cp]), expression(hat(F)[bic])))
par(op)
```

mable.dr

MABLE in Density Ratio Model

Description

Maximum approximate Bernstein/Beta likelihood estimation in a density ratio model based on two-sample raw data.

Usage

```
mable.dr(
  x,
  y,
  M,
  regr,
  ...,
  interval = c(0, 1),
  alpha = NULL,
  vb = 0,
```

```

baseline = NULL,
controls = mable.ctrl(),
progress = TRUE,
message = FALSE
)

```

Arguments

<code>x, y</code>	original two sample raw data, <code>x:"Control", y: "Case"</code> .
<code>M</code>	a positive integer or a vector (<code>m0, m1</code>).
<code>regr</code>	regressor vector function $r(x) = (1, r_1(x), \dots, r_d(x))$ which returns $n \times (d+1)$ matrix, $n=\text{length}(x)$
<code>...</code>	additional arguments to be passed to <code>regr</code>
<code>interval</code>	a vector (<code>a, b</code>) containing the endpoints of supporting/truncation interval of <code>x</code> and <code>y</code> .
<code>alpha</code>	initial regression coefficient, missing value is imputed by logistic regression
<code>vb</code>	code for vanishing boundary constraints, -1: $f_0(a)=0$ only, 1: $f_0(b)=0$ only, 2: both, 0: none (default).
<code>baseline</code>	the working baseline, "Control" or "Case", if NULL it is chosen to the one with smaller estimated lower bound for model degree.
<code>controls</code>	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion for EM and Newton iterations. Default is <code>mable.ctrl</code> . See Details.
<code>progress</code>	logical: should a text progressbar be displayed
<code>message</code>	logical: should warning messages be displayed

Details

Suppose that `x` ("control") and `y` ("case") are independent samples from f_0 and f_1 which samples satisfy $f_1(x)=f_0(x)\exp[\alpha_0+\alpha'r(x)]$ with $r(x)=(r_1(x), \dots, r_d(x))$. Maximum approximate Bernstein/Beta likelihood estimates of (α_0, α) , f_0 and f_1 are calculated. If support is (a, b) then replace $r(x)$ by $r[a+(b-a)x]$. For a fixed m , using the Bernstein polynomial model for baseline f_0 , MABLEs of f_0 and parameters α can be estimated by EM algorithm and Newton iteration. If estimated lower bound m_b for m based on `y` is smaller than that based on `x`, then switch `x` and `y` and f_1 is used as baseline. If $M=m$ or $m_0=m_1=m$, then m is a preselected degree. If $m_0 < m_1$ it specifies the set of consecutive candidate model degrees $m_0:m_1$ for searching an optimal degree by the change-point method, where $m_1 - m_0 > 3$.

Value

A list with components

- `m` the given or a selected degree by method of change-point
- `p` the estimated vector of mixture proportions $p = (p_0, \dots, p_m)$ with the given or selected degree `m`
- `alpha` the estimated regression coefficients
- `mloglik` the maximum log-likelihood at degree `m`

- interval support/truncation interval (a,b)
- baseline ="control" if f_0 is used as baseline, or ="case" if f_1 is used as baseline.
- M the vector (m0, m1), where m1, if greater than m0, is the largest candidate when the search stopped
- lk log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- lr likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- pval the p-values of the change-point tests for choosing optimal model degree
- chpts the change-points chosen with the given candidate model degrees

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z., Maximum Approximate Bernstein Likelihood Estimation of Densities in a Two-sample Semiparametric Model

Examples

```
# Hosmer and Lemeshow (1989):
# ages and the status of coronary disease (CHD) of 100 subjects
x<-c(20, 23, 24, 25, 26, 26, 28, 28, 29, 30, 30, 30, 30, 30, 32,
32, 33, 33, 34, 34, 34, 34, 34, 35, 35, 36, 36, 37, 37, 38, 38, 39,
40, 41, 41, 42, 42, 42, 43, 43, 44, 44, 45, 46, 47, 47, 48, 49,
49, 50, 51, 52, 55, 57, 57, 58, 60, 64)
y<-c(25, 30, 34, 36, 37, 39, 40, 42, 43, 44, 44, 45, 46, 47, 48,
48, 49, 50, 52, 53, 53, 54, 55, 55, 56, 56, 56, 57, 57, 57, 57,
58, 58, 59, 59, 60, 61, 62, 62, 63, 64, 65, 69)
regr<-function(x) cbind(1,x)
chd.mable<-mable.dr(x, y, M=c(1, 15), regr, interval = c(20, 70))
chd.mable
```

mable.dr.group

Mable fit of the density ratio model based on grouped data

Description

Maximum approximate Bernstein/Beta likelihood estimation in a density ratio model based on two-sample grouped data.

Usage

```
mable.dr.group(
  t,
  n0,
  n1,
  M,
  regr,
  ...,
  interval = c(0, 1),
  alpha = NULL,
  vb = 0,
  controls = mable.ctrl(),
  progress = TRUE,
  message = TRUE
)
```

Arguments

t	cutpoints of class intervals
n0, n1	frequencies of two sample data grouped by the classes specified by t. n0: "Control", n1: "Case".
M	a positive integer or a vector (m0, m1).
regr	regressor vector function $r(x) = (1, r_1(x), \dots, r_d(x))$ which returns n x (d+1) matrix, n=length(x)
...	additional arguments to be passed to regr
interval	a vector (a,b) containing the endpoints of supporting/truncation interval of x and y.
alpha	a given regression coefficient, missing value is imputed by logistic regression
vb	code for vanishing boundary constraints, -1: f0(a)=0 only, 1: f0(b)=0 only, 2: both, 0: none (default).
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion for EM and Newton iterations. Default is <code>mable.ctrl</code> . See Details.
progress	logical: should a text progressbar be displayed
message	logical: should warning messages be displayed

Details

Suppose that n_0 ("control") and n_1 ("case") are frequencies of independent samples grouped by the classes t from f_0 and f_1 which satisfy $f_1(x) = f_0(x) \exp[\alpha_0 + \alpha' r(x)]$ with $r(x) = (r_1(x), \dots, r_d(x))$. Maximum approximate Bernstein/Beta likelihood estimates of (α_0, α) , f_0 and f_1 are calculated. If support is (a, b) then replace $r(x)$ by $r[a + (b-a)x]$. For a fixed m , using the Bernstein polynomial model for baseline f_0 , MABLEs of f_0 and parameters α can be estimated by EM algorithm and Newton iteration. If estimated lower bound m_b for m based on n_1 is smaller than that based on n_0 , then switch n_0 and n_1 and use f_1 as baseline. If $M = m$ or $m_0 = m_1 = m$, then m is a preselected degree. If $m_0 < m_1$ it specifies the set of consecutive candidate model degrees $m_0 : m_1$ for searching an optimal degree by the change-point method, where $m_1 - m_0 > 3$.

mable.group	<i>Mable fit of one-sample grouped data by an optimal or a preselected model degree</i>
-------------	---

Description

Maximum approximate Bernstein/Beta likelihood estimation based on one-sample grouped data with an optimal selected by the change-point method among $m_0:m_1$ or a preselected model degree m .

Usage

```
mable.group(
  x,
  breaks,
  M,
  interval = c(0, 1),
  IC = c("none", "aic", "hqic", "all"),
  vb = 0,
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

x	vector of frequencies
breaks	class interval end points
M	a positive integer or a vector (m_0, m_1). If $M = m$ or $m_0 = m_1 = m$, then m is a preselected degree. If $m_0 < m_1$ it specifies the set of consecutive candidate model degrees $m_0:m_1$ for searching an optimal degree, where $m_1 - m_0 > 3$.
interval	a vector containing the endpoints of support/truncation interval
IC	information criterion(s) in addition to Bayesian information criterion (BIC). Current choices are "aic" (Akaike information criterion) and/or "hqic" (Hannan–Quinn information criterion).
vb	code for vanishing boundary constraints, -1: $f_0(a)=0$ only, 1: $f_0(b)=0$ only, 2: both, 0: none (default).
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion eps. Default is <code>mable.ctrl</code> . See Details.
progress	if TRUE a text progressbar is displayed

Details

Any continuous density function f on a known closed supporting interval $[a, b]$ can be estimated by Bernstein polynomial $f_m(x; p) = \sum_{i=0}^m p_i \beta_{mi}[(x-a)/(b-a)]/(b-a)$, where $p = (p_0, \dots, p_m)$, $p_i \geq 0$, $\sum_{i=0}^m p_i = 1$ and $\beta_{mi}(u) = (m+1) \binom{m}{i} u^i (1-u)^{m-i}$, $i = 0, 1, \dots, m$, is the beta density

with shapes $(i+1, m-i+1)$. For each m , the MABLE of the coefficients p , the mixture proportions, are obtained using EM algorithm. The EM iteration for each candidate m stops if either the total absolute change of the log likelihood and the coefficients of Bernstein polynomial is smaller than `eps` or the maximum number of iterations `maxit` is reached.

If $m_0 < m_1$, an optimal model degree is selected as the change-point of the increments of log-likelihood, log likelihood ratios, for $m \in \{m_0, m_0 + 1, \dots, m_1\}$. Alternatively, one can choose an optimal degree based on the BIC (Schwarz, 1978) which are evaluated at $m \in \{m_0, m_0 + 1, \dots, m_1\}$. The search for optimal degree m is stopped if either m_1 is reached with a warning or the test for change-point results in a p-value `pval` smaller than `sig.level`. The BIC for a given degree m is calculated as in Schwarz (1978) where the dimension of the model is $d = \#\{i : \hat{p}_i \geq \epsilon, i = 0, \dots, m\} - 1$ and a default ϵ is chosen as `.Machine$double.eps`.

Value

A list with components

- `m` the given or a selected degree by method of change-point
- `p` the estimated p with degree m
- `mloglik` the maximum log-likelihood at degree m
- `interval` supporting interval (a, b)
- `convergence` An integer code. 0 indicates successful completion (all the EM iterations are convergent and an optimal degree is successfully selected in `M`). Possible error codes are
 - 1, indicates that the iteration limit `maxit` had been reached in at least one EM iteration;
 - 2, the search did not finish before `m1`.
- `delta` the convergence criterion `delta` value

and, if $m_0 < m_1$,

- `M` the vector (m_0, m_1) , where m_1 , if greater than m_0 , is the largest candidate when the search stopped
- `lk` log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- `lr` likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- `ic` a list containing the selected information criterion(s)
- `pval` the p-values of the change-point tests for choosing optimal model degree
- `chpts` the change-points chosen with the given candidate model degrees

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. (2017) Bernstein polynomial model for grouped continuous data. *Journal of Nonparametric Statistics*, 29(4):831-848.

See Also[mable.ic](#)**Examples**

```
## Chicken Embryo Data
data(chicken.embryo)
a<-0; b<-21
day<-chicken.embryo$day
nT<-chicken.embryo$nT
Day<-rep(day,nT)
res<-mable.group(x=nT, breaks=a:b, M=c(2,100), interval=c(a, b), IC="aic",
  controls=mable.ctrl(sig.level=1e-6, maxit=2000, eps=1.0e-7))
op<-par(mfrow=c(1,2), lwd=2)
layout(rbind(c(1, 2), c(3, 3)))
plot(res, which="likelihood")
plot(res, which="change-point")
fk<-density(x=rep((0:20)+.5, nT), bw="sj", n=101, from=a, to=b)
hist(Day, breaks=seq(a,b, length=12), freq=FALSE, col="grey",
  border="white", main="Histogram and Density Estimates")
plot(res, which="density", types=1:2, colors=1:2)
lines(fk, lty=2, col=2)
legend("topright", lty=c(1:2), c("MABLE", "Kernel"), bty="n", col=c(1:2))
par(op)
```

mable.hellcorr	<i>Estimate of Hellinger Correlation between two random variables and Bootstrap</i>
----------------	---

Description

Estimate of Hellinger Correlation between two random variables and Bootstrap

Usage

```
mable.hellcorr(
  x,
  unif.mar = FALSE,
  pseudo.obs = c("empirical", "mable"),
  M0 = c(1, 1),
  M = c(30, 30),
  search = TRUE,
  mar.deg = TRUE,
  high.dim = FALSE,
  interval = cbind(0:1, 0:1),
  B = 200L,
  conf.level = 0.95,
```

```

    integral = TRUE,
    controls = mable.ctrl(sig.level = 0.05),
    progress = FALSE
  )

hellcorr(
  x,
  unif.mar = FALSE,
  pseudo.obs = c("empirical", "mable"),
  M0 = c(1, 1),
  M = c(30, 30),
  search = TRUE,
  mar.deg = TRUE,
  high.dim = FALSE,
  interval = cbind(0:1, 0:1),
  B = 200L,
  conf.level = 0.95,
  integral = TRUE,
  controls = mable.ctrl(sig.level = 0.05),
  progress = FALSE
)

```

Arguments

<code>x</code>	an $n \times 2$ data matrix of observations of the two random variables
<code>unif.mar</code>	logical, whether all the marginals distributions are uniform or not. If not the pseudo observations will be created using empirical or mable marginal distributions.
<code>pseudo.obs</code>	"empirical": use empirical distribution to form pseudo, observations, or "mable": use mable of marginal cdfs to form pseudo observations
<code>M0</code>	a nonnegative integer or a vector of d nonnegative integers specify starting candidate degrees for searching optimal degrees.
<code>M</code>	a positive integer or a vector of d positive integers specify the maximum candidate or the given model degrees for the joint density.
<code>search</code>	logical, whether to search optimal degrees between <code>M0</code> and <code>M</code> or not but use <code>M</code> as the given model degrees for the joint density.
<code>mar.deg</code>	logical, if TRUE (default), the optimal degrees are selected based on marginal data, otherwise, the optimal degrees are chosen by the method of change-point. See details.
<code>high.dim</code>	logical, data are high dimensional/large sample or not if TRUE, run a slower version procedure which requires less memory
<code>interval</code>	a 2 by 2 matrix, columns are the marginal supports
<code>B</code>	the number of bootstrap samples and number of Monte Carlo runs for estimating p.value of the test for Hellinger correlation = 0 if test=TRUE.
<code>conf.level</code>	confidence level
<code>integral</code>	logical, using "integrate()" or not (Riemann sum)

controls Object of class `mable.ctrl()` specifying iteration limit and the convergence criterion `eps`. Default is `mable.ctrl`. See Details.

progress if TRUE a text progressbar is displayed

Details

This function calls `mable.copula()` for estimation of the copula density.

Value

- eta Hellinger correlation
- CI.eta Bootstrap confidence interval for Hellinger correlation if $B > 0$.

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z., Nonparametric Maximum Likelihood Estimation of Copula

See Also

[mable](#), [mable.mvar](#), [mable.copula](#)

mable.ic

Mable fit based on one-sample interval censored data

Description

Maximum approximate Bernstein/Beta likelihood estimation of density and cumulative/survival distributions functions based on interval censored event time data.

Usage

```
mable.ic(  
  data,  
  M,  
  pi0 = NULL,  
  tau = Inf,  
  IC = c("none", "aic", "hqic", "all"),  
  controls = mable.ctrl(),  
  progress = TRUE  
)
```

Arguments

data	a dataset either data.frame or an $n \times 2$ matrix.
M	an positive integer or a vector (m_0, m_1) . If $M = m$ or $m_0 = m_1 = m$, then m is a preselected degree. If $m_0 < m_1$ it specifies the set of consecutive candidate model degrees $m_0:m_1$ for searching an optimal degree, where $m_1 - m_0 > 3$.
pi0	Initial guess of $\pi = F(\tau_n)$. Without right censored data, $pi0 = 1$. See 'Details'.
tau	right endpoint of support $[0, \tau)$ must be greater than or equal to the maximum observed time
IC	information criterion(s) in addition to Bayesian information criterion (BIC). Current choices are "aic" (Akaike information criterion) and/or "qhic" (Hannan-Quinn information criterion).
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

Let $f(t)$ and $F(t) = 1 - S(t)$ be the density and cumulative distribution functions of the event time, respectively. Then $f(t)$ on $[0, \tau_n]$ can be approximated by $f_m(t; p) = \tau_n^{-1} \sum_{i=0}^m p_i \beta_{mi}(t/\tau_n)$, where $p_i \geq 0$, $i = 0, \dots, m$, $\sum_{i=0}^m p_i = 1 - p_{m+1}$, $\beta_{mi}(u)$ is the beta density with shapes $i + 1$ and $m - i + 1$, and τ_n is the largest observed time, either uncensored time, or right endpoint of interval/left censored, or left endpoint of right censored time. We can approximate $S(t)$ on $[0, \tau]$ by $S_m(t; p) = \sum_{i=0}^{m+1} p_i \bar{B}_{mi}(t/\tau)$, where $\bar{B}_{mi}(u)$, $i = 0, \dots, m$, is the beta survival function with shapes $i + 1$ and $m - i + 1$, $\bar{B}_{m,m+1}(t) = 1$, $p_{m+1} = 1 - \pi$, and $\pi = F(\tau_n)$. For data without right-censored time, $p_{m+1} = 1 - \pi = 0$. The search for optimal degree m is stopped if either m_1 is reached or the test for change-point results in a p-value `pval` smaller than `sig.level`.

Each row of data, (l, u) , is the interval containing the event time. Data is uncensored if $l = u$, right censored if $u = \text{Inf}$ or $u = \text{NA}$, and left censored data if $l = 0$.

Value

a class 'mable' object with components

- `p` the estimated p with degree m selected by the change-point method
- `mloglik` the maximum log-likelihood at an optimal degree m
- `interval` support/truncation interval (θ, b)
- `M` the vector (m_0, m_1) , where m_1 is the last candidate when the search stopped
- `m` the selected optimal degree by the method of change-point
- `lk` log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- `lr` likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- `tau.n` maximum observed time τ_n
- `tau` right endpoint of support $[0, \tau)$
- `ic` a list containing the selected information criterion(s)

- pval the p-values of the change-point tests for choosing optimal model degree
- chpts the change-points chosen with the given candidate model degrees
- convergence an integer code. 0 indicates successful completion(the iteration is convergent). 1 indicates that the maximum candidate degree had been reached in the calculation;
- delta the final pval of the change-point for selecting the optimal degree m;

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. (2019) Maximum Approximate Bernstein Likelihood Estimation in Proportional Hazard Model for Interval-Censored Data, arXiv:1906.08882 .

See Also

[mable.group](#)

Examples

```
library(mable)
bcos=cosmesis
bc.res0<-mable.ic(bcos[bcos$treat=="RT",1:2], M=c(1,50), IC="none")
bc.res1<-mable.ic(bcos[bcos$treat=="RCT",1:2], M=c(1,50), IC="none")
op<-par(mfrow=c(2,2),lwd=2)
plot(bc.res0, which="change-point", lgd.x="right")
plot(bc.res1, which="change-point", lgd.x="right")
plot(bc.res0, which="survival", add=FALSE, xlab="Months", ylim=c(0,1), main="Radiation Only")
legend("topright", bty="n", lty=1:2, col=1:2, c(expression(hat(S)[CP]),
expression(hat(S)[BIC])))
plot(bc.res1, which="survival", add=FALSE, xlab="Months", main="Radiation and Chemotherapy")
legend("topright", bty="n", lty=1:2, col=1:2, c(expression(hat(S)[CP]),
expression(hat(S)[BIC])))
par(op)
```

mable.mvar

*Maximum Approximate Bernstein Likelihood Estimate of Multivariate
Density Function*

Description

Maximum Approximate Bernstein Likelihood Estimate of Multivariate Density Function

Usage

```
mable.mvar(
  x,
  M0 = 1L,
  M,
  search = TRUE,
  interval = NULL,
  mar.deg = TRUE,
  method = c("cd", "em", "lmem"),
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

x	an n x d matrix or data.frame of multivariate sample of size n
M0	a positive integer or a vector of d positive integers specify starting candidate degrees for searching optimal degrees.
M	a positive integer or a vector of d positive integers specify the maximum candidate or the given model degrees for the joint density.
search	logical, whether to search optimal degrees between M0 and M or not but use M as the given model degrees for the joint density.
interval	a vector of two endpoints or a 2 x d matrix, each column containing the endpoints of support/truncation interval for each marginal density. If missing, the i-th column is assigned as $c(\min(x[, i]), \max(x[, i]))$.
mar.deg	logical, if TRUE, the optimal degrees are selected based on marginal data, otherwise, the optimal degrees are chosen the joint data. See details.
method	method for finding maximum likelihood estimate. "cd": coordinate-descent; less memory for data that are high dimensional/large sample.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion eps. Default is <code>mable.ctrl</code> . See Details.
progress	if TRUE a text progressbar is displayed

Details

A d -variate density f on a hyperrectangle $[a, b] = [a_1, b_1] \times \dots \times [a_d, b_d]$ can be approximated by a mixture of d -variate beta densities on $[a, b]$, $\beta_{m_j}(x) = \prod_{i=1}^d \beta_{m_i, j_i}[(x_i - a_i)/(b_i - a_i)]/(b_i - a_i)$, with proportion $p(j_1, \dots, j_d)$, $0 \leq j_i \leq m_i$, $i = 1, \dots, d$. If `search=TRUE` then the model degrees are chosen using a method of change-point based on the marginal data if `mar.deg=TRUE` or the joint data if `mar.deg=FALSE`. If `search=FALSE`, then the model degree is specified by M . For large data and multimodal density, the search for the model degrees is very time-consuming. In this case, it is suggested that use `method="cd"` and select the degrees based on marginal data using `mable` or `optimable`.

Value

A list with components

- `m` a vector of the selected optimal degrees by the method of change-point
- `p` a vector of the mixture proportions $p(j_1, \dots, j_d)$, arranged in the column-major order of $j = (j_1, \dots, j_d)$, $0 \leq j_i \leq m_i, i = 1, \dots, d$.
- `mloglik` the maximum log-likelihood at an optimal degree `m`
- `pval` the p-values of change-points for choosing the optimal degrees for the marginal densities
- `M` the vector (m_1, m_2, \dots, m_d) , where m_i is the largest candidate degree when the search stopped for the i -th marginal density
- `interval` support hyperrectangle $[a, b] = [a_1, b_1] \times \dots \times [a_d, b_d]$
- `convergence` An integer code. 0 indicates successful completion (the EM iteration is convergent). 1 indicates that the iteration limit `maxit` had been reached in the EM iteration;

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. (2016) Efficient and robust density estimation using Bernstein type polynomials. *Journal of Nonparametric Statistics*, 28(2):250-271. Wang, T. and Guan, Z., (2019) Bernstein Polynomial Model for Nonparametric Multivariate Density, *Statistics*, Vol. 53, no. 2, 321-338

See Also

[mable](#), [optimable](#)

Examples

```
## Old Faithful Data
a<-c(0, 40); b<-c(7, 110)
ans<- mable.mvar(faithful, M = c(46,19), search =FALSE, method="em",
  interval = rbind(a,b), progress=FALSE)
plot(ans, which="density")
plot(ans, which="cumulative")
```

mable.ph

Mable fit of Cox's proportional hazards regression model

Description

Maximum approximate Bernstein/Beta likelihood estimation in Cox's proportional hazards regression model based on interval censored event time data.

Usage

```
mable.ph(
  formula,
  data,
  M,
  g = NULL,
  p = NULL,
  pi0 = NULL,
  tau = Inf,
  x0 = NULL,
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

formula	regression formula. Response must be cbind. See 'Details'.
data	a data frame containing variables in formula.
M	a positive integer or a vector (m0, m1). If M = m or m0 = m1, then m0 is a preselected degree. If m0 < m1 it specifies the set of consecutive candidate model degrees m0:m1 for searching an optimal degree, where m1 - m0 > 3.
g	initial guess of d -vector of regression coefficients. See 'Details'.
p	an initial coefficients of Bernstein polynomial model of degree m0, default is the uniform initial.
pi0	Initial guess of $\pi(x_0) = F(\tau_n x_0)$. Without right censored data, pi0 = 1. See 'Details'.
tau	right endpoint of support $[0, \tau)$ must be greater than or equal to the maximum observed time
x0	a data frame specifying working baseline covariates on the right-hand-side of formula. See 'Details'.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

Consider Cox's PH model with covariate for interval-censored failure time data: $S(t|x) = S(t|x_0)^{\exp(\gamma^T(x-x_0))}$, where x_0 satisfies $\gamma^T(x-x_0) \geq 0$, where x and x_0 may contain dummy variables and interaction terms. The working baseline x_0 in arguments contains only the values of terms excluding dummy variables and interaction terms in the right-hand-side of formula. Thus g is the initial guess of the coefficients γ of $x-x_0$ and could be longer than x_0 . Let $f(t|x)$ and $F(t|x) = 1 - S(t|x)$ be the density and cumulative distribution functions of the event time given $X = x$, respectively. Then $f(t|x_0)$ on $[0, \tau_n]$ can be approximated by $f_m(t|x_0, p) = \tau_n^{-1} \sum_{i=0}^m p_i \beta_{mi}(t/\tau_n)$, where $p_i \geq 0$, $i = 0, \dots, m$, $\sum_{i=0}^m p_i = 1 - p_{m+1}$, $\beta_{mi}(u)$ is the beta density with shapes $i+1$ and $m-i+1$, and τ_n is the largest observed time, either uncensored time, or right endpoint of interval/left censored, or left endpoint of right censored time. So we can approximate $S(t|x_0)$ on $[0, \tau_n]$ by $S_m(t|x_0; p) = \sum_{i=0}^{m+1} p_i \bar{B}_{mi}(t/\tau_n)$, where $\bar{B}_{mi}(u)$, $i = 0, \dots, m$, is the beta survival function with shapes $i+1$ and $m-i+1$, $\bar{B}_{m,m+1}(t) = 1$, $p_{m+1} = 1 - \pi(x_0)$, and $\pi(x_0) = F(\tau_n|x_0)$. For data without right-censored time, $p_{m+1} = 1 - \pi(x_0) = 0$.

Response variable should be of the form `cbind(1, u)`, where $(1, u)$ is the interval containing the event time. Data is uncensored if $1 = u$, right censored if $u = \text{Inf}$ or $u = \text{NA}$, and left censored data if $1 = 0$. The associated covariate contains d columns. The baseline x_0 should be chosen so that $\gamma'(x-x_0)$ is nonnegative for all the observed x and all γ in a neighborhood of its true value.

A missing initial value of g is imputed by `ic_sp()` of package `icenReg`.

The search for optimal degree m stops if either `m1` is reached or the test for change-point results in a p -value `pval` smaller than `sig.level`. This process takes longer than `maple.ph` to select an optimal degree.

Value

A list with components

- `m` the selected/preselected optimal degree m
- `p` the estimate of $p = (p_0, \dots, p_m, p_{m+1})$, the coefficients of Bernstein polynomial of degree m
- `coefficients` the estimated regression coefficients of the PH model
- `SE` the standard errors of the estimated regression coefficients
- `z` the z-scores of the estimated regression coefficients
- `mloglik` the maximum log-likelihood at an optimal degree m
- `tau.n` maximum observed time τ_n
- `tau` right endpoint of support $[0, \tau)$
- `x0` the working baseline covariates
- `egx0` the value of $e^{\gamma'x_0}$
- `convergence` an integer code, 1 indicates either the EM-like iteration for finding maximum likelihood reached the maximum iteration for at least one m or the search of an optimal degree using change-point method reached the maximum candidate degree, 2 indicates both occurred, and 0 indicates a successful completion.
- `delta` the final `delta` if `m0 = m1` or the final `pval` of the change-point for searching the optimal degree m ;

and, if $m_0 < m_1$,

- M the vector (m_0, m_1) , where m_1 is the last candidate degree when the search stopped
- lk log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- lr likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- $pval$ the p-values of the change-point tests for choosing optimal model degree
- $chpts$ the change-points chosen with the given candidate model degrees

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. Maximum Approximate Bernstein Likelihood Estimation in Proportional Hazard Model for Interval-Censored Data, *Statistics in Medicine*. 2020; 1–21. <https://doi.org/10.1002/sim.8801>.

See Also

[maple.ph](#)

Examples

```
# Ovarian Cancer Survival Data
require(survival)
fuptime2<-ovarian$fuptime
fuptime2[ovarian$fustat==0]<-Inf
ovarian2<-data.frame(age=ovarian$age, fuptime1=ovarian$fuptime,
  fuptime2=fuptime2)
ova<-mable.ph(cbind(fuptime1, fuptime2) ~ age, data = ovarian2,
  M=c(2,35), g=.16, x0=data.frame(age=35))
op<-par(mfrow=c(2,2))
plot(ova, which = "likelihood")
plot(ova, which = "change-point")
plot(ova, y=data.frame(age=60), which="survival", add=FALSE, type="l",
  xlab="Days", main="Age = 60")
plot(ova, y=data.frame(age=65), which="survival", add=FALSE, type="l",
  xlab="Days", main="Age = 65")
par(op)
```

mable.po

Mable fit of proportional odds rate regression model

Description

Maximum approximate Bernstein/Beta likelihood estimation in general proportional odds regression model based on interval censored event time data.

Usage

```
mable.po(
  formula,
  data,
  M,
  g = NULL,
  tau,
  x0 = NULL,
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

formula	regression formula. Response must be cbind. See 'Details'.
data	a data frame containing variables in formula.
M	a positive integer or a vector (m0, m1). If M = m or m0 = m1 = m, then m is a preselected degree. If m0 < m1 it specifies the set of consecutive candidate model degrees m0:m1 for searching an optimal degree, where m1 - m0 > 3.
g	an initial guess of d -vector of regression coefficients. See 'Details'.
tau	right endpoint of support $[0, \tau]$ must be greater than or equal to the maximum observed time
x0	a data frame specifying working baseline covariates on the right-hand-side of formula. See 'Details'.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

Consider PO model with covariate for interval-censored failure time data: $[1 - S(t|x)]/S(t|x) = \exp[\gamma'(x - x_0)][1 - S(t|x_0)]/S(t|x_0)$, where x_0 satisfies $\gamma'(x - x_0) \geq 0$, where x and x_0 may contain dummy variables and interaction terms. The working baseline x_0 in arguments contains only the values of terms excluding dummy variables and interaction terms in the right-hand-side of formula. Thus g is the initial guess of the coefficients γ of $x - x_0$ and could be longer than x_0 . Let $f(t|x)$ and $F(t|x) = 1 - S(t|x)$ be the density and cumulative distribution functions of the event time given $X = x$, respectively. Then $f(t|x_0)$ on $[0, \tau]$ can be approximated by $f_m(t|x_0; p) = \tau^{-1} \sum_{i=0}^m p_i \beta_{mi}(t/\tau)$, where $p_i \geq 0$, $i = 0, \dots, m$, $\sum_{i=0}^m p_i = 1$, $\beta_{mi}(u)$ is the beta density with shapes $i + 1$ and $m - i + 1$, and τ is the right endpoint of support interval of the baseline density. We can approximate $S(t|x_0)$ on $[0, \tau]$ by $S_m(t|x_0; p) = \sum_{i=0}^m p_i \bar{B}_{mi}(t/\tau)$, where $\bar{B}_{mi}(u)$, $i = 0, \dots, m$, is the beta survival function with shapes $i + 1$ and $m - i + 1$.

Response variable should be of the form `cbind(1, u)`, where $(1, u)$ is the interval containing the event time. Data is uncensored if $1 = u$, right censored if $u = \text{Inf}$ or $u = \text{NA}$, and left censored if $1 = 0$. The associated covariate contains d columns. The baseline x_0 should be chosen so that $\gamma'(x - x_0)$ is nonnegative for all the observed x and all γ in a neighborhood of its true value.

A missing initial value of g is imputed by `ic_sp()` of package `icenReg` with `model="po"`. The search for optimal degree m stops if either m_1 is reached or the test for change-point results in a p -value `pval` smaller than `sig.level`. This process takes longer than `maple.po` to select an optimal degree.

Value

A list with components

- `m` the selected/preselected optimal degree m
- `p` the estimate of $p = (p_0, \dots, p_m)$, the coefficients of Bernstein polynomial of degree m
- `coefficients` the estimated regression coefficients of the PO model
- `SE` the standard errors of the estimated regression coefficients
- `z` the z -scores of the estimated regression coefficients
- `mloglik` the maximum log-likelihood at an optimal degree m
- `tau.n` maximum observed time τ_n
- `tau` right endpoint of support $[0, \tau]$
- `x0` the working baseline covariates
- `egx0` the value of $e^{\gamma'x_0}$
- `convergence` an integer code, 1 indicates either the EM-like iteration for finding maximum likelihood reached the maximum iteration for at least one m or the search of an optimal degree using change-point method reached the maximum candidate degree, 2 indicates both occurred, and 0 indicates a successful completion.
- `delta` the final `delta` if `m0 = m1` or the final `pval` of the change-point for searching the optimal degree m ;

and, if `m0 < m1`,

- `M` the vector (m_0, m_1) , where m_1 is the last candidate degree when the search stopped
- `lk` log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- `lr` likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- `pval` the p -values of the change-point tests for choosing optimal model degree
- `chpts` the change-points chosen with the given candidate model degrees

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. Maximum Likelihood Estimation in Proportional Odds Regression Model Based on Interval-Censored Event-time Data

See Also

[maple.ph](#)

Examples

```

# Veteran's Administration Lung Cancer Data
require(survival)
require(icenReg)
require(mable)
l<-veteran$time->u
u[veteran$status==0]<-Inf
veteran1<-data.frame(l=l, u=u, karno=veteran$karno, celltype=veteran$celltype,
  trt=veteran$trt, age=veteran$age, prior=veteran$prior>0)
fit.sp<-ic_sp(cbind(l,u) ~ karno+celltype, data = veteran1, model="po")
x0<-data.frame(karno=100, celltype="squamous")
tau<-2000
res<-mable.po(cbind(l,u) ~ karno+celltype, data = veteran1, M=c(1,35),
  g=-fit.sp$coefficients, x0=x0, tau=tau)
op<-par(mfrow=c(2,2))
plot(res, which = "likelihood")
plot(res, which = "change-point")
plot(res, y=data.frame(karno=20, celltype="squamous"), which="survival",
  add=FALSE, type="l", xlab="Days",
  main=expression(paste("Survival: ", bold(x)==0)))
plot(res, y=data.frame(karno=80, celltype="smallcell"), which="survival",
  add=FALSE, type="l", xlab="Days",
  main=expression(paste("Survival: ", bold(x)==bold(x)[0])))
par(op)

```

mable.reg

Mable fit of semiparametric regression model based on interval censored data

Description

Wrapping all mable fit of regression models in one function. Using maximum approximate Bernstein/Beta likelihood estimation to fit semiparametric regression models: Cox ph model, proportional odds(po) model, accelerated failure time model, and so on.

Usage

```

mable.reg(
  formula,
  data,
  model = c("ph", "aft", "po"),
  M,
  g = NULL,
  pi0 = NULL,
  tau = Inf,
  x0 = NULL,
  controls = mable.ctrl(),

```

```

    progress = TRUE
  )

```

Arguments

formula	regression formula. Response must be of the form <code>cbind(1, u)</code> . See 'Details'.
data	a data frame containing variables in formula.
model	the model to fit. Current options are "ph" (Cox PH) or "aft" (accelerated failure time model)
M	a vector (m_0, m_1) specifies the set of consecutive integers as candidate degrees
g	an initial guess of the regression coefficients
pi0	Initial guess of $\pi(x_0) = F(\tau_n x_0)$. Without right censored data, $\text{pi}0 = 1$. See 'Details'.
tau	right endpoint of support $[0, \tau)$ must be greater than or equal to the maximum observed time
x0	a data frame containing working baseline covariates on the right-hand-side of formula. See 'Details'.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

For "ph" model a missing initial guess of the regression coefficients `g` is obtained by `ic_sp()` of package `icenReg`. For "aft" model a missing `g` is imputed by the rank estimate `aftsrr()` of package `aftgee` for right-censored data. For general interval censored observations, we keep the right-censored but replace the finite interval with its midpoint and fit the data by `aftsrr()` as a right-censored data.

Value

A 'mable_reg' class object

Author(s)

Zhong Guan <zguan@iu.edu>

See Also

[mable.aft](#), [mable.ph](#), [mable.po](#)

made.copula *Minimum Approximate Distance Estimate of Copula Density*

Description

Minimum Approximate Distance Estimate of Copula Density

Usage

```
made.copula(
  x,
  unif.mar = FALSE,
  M = 30,
  search = TRUE,
  interval = NULL,
  pseudo.obs = c("empirical", "mable"),
  sig.level = 0.01
)
```

Arguments

x	an n x d matrix of data values
unif.mar	marginals are all uniform (x contain pseudo observations) or not.
M	d-vector of preselected or maximum model degrees
search	logical, whether to search optimal degrees between 0 and M or not but use M as the given model degrees for the joint density.
interval	a 2 by d matrix specifying the support/truncate interval of x, if unif.mar=TRUE then interval is the unit hypercube
pseudo.obs	When unif.mar=FALSE, use "empirical" distribution to create pseudo observations, or use "mable" of marginal cdfs to create pseudo observations
sig.level	significance level for p-value of change-point

Details

With given model degrees m , the parameters p , the mixing proportions of the beta distribution, are calculated as the minimizer of the approximate L_2 distance between the empirical distribution and the Bernstein polynomial model. The optimal model degrees m are chosen by a change-point method. The quadratic programming with linear constraints is used to solve the problem.

Value

An invisible mable object with components

- m the given degree
- p the estimated vector of mixture proportions $p = (p_0, \dots, p_m)$ with the given degree m
- D the minimum distance at degree m

made.density	<i>Minimum Approximate Distance Estimate of Density Function with an optimal model degree</i>
--------------	---

Description

Minimum Approximate Distance Estimate of Density Function with an optimal model degree

Usage

```
made.density(
  x,
  M0 = 1L,
  M,
  search = TRUE,
  interval = NULL,
  mar.deg = TRUE,
  method = c("qp", "em"),
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

x	an n x d matrix or data.frame of multivariate sample of size n
M0	a positive integer or a vector of d positive integers specify starting candidate degrees for searching optimal degrees.
M	a positive integer or a vector of d positive integers specify the maximum candidate or the given model degrees for the joint density.
search	logical, whether to search optimal degrees between M0 and M or not but use M as the given model degrees for the joint density.
interval	a vector of two endpoints or a 2 x d matrix, each column containing the endpoints of support/truncation interval for each marginal density. If missing, the i-th column is assigned as $c(\min(x[, i]), \max(x[, i]))$.
mar.deg	logical, if TRUE, the optimal degrees are selected based on marginal data, otherwise, the optimal degrees are chosen the joint data. See details.
method	method for finding minimum distance estimate. "em": EM like method;
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion eps. Default is <code>mable.ctrl</code> . See Details.
progress	if TRUE a text progressbar is displayed

Details

A d -variate cdf F on a hyperrectangle $[a, b] = [a_1, b_1] \times \cdots \times [a_d, b_d]$ can be approximated by a mixture of d -variate beta cdfs on $[a, b]$, $\beta_{mj}(x) = \prod_{i=1}^d B_{m_i, j_i}[(x_i - a_i)/(b_i - a_i)]$, with proportion $p(j_1, \dots, j_d)$, $0 \leq j_i \leq m_i, i = 1, \dots, d$. With a given model degree m , the parameters p , the mixing proportions of the beta distribution, are calculated as the minimizer of the approximate L_2 distance between the empirical distribution and the Bernstein polynomial model. The quadratic programming with linear constraints is used to solve the problem. If `search=TRUE` then the model degrees are chosen using a method of change-point based on the marginal data if `mar.deg=TRUE` or the joint data if `mar.deg=FALSE`. If `search=FALSE`, then the model degree is specified by M .

Value

An invisible mable object with components

- `m` the given model degree(s)
- `p` the estimated vector of mixture proportions with the given optimal degree(s) `m`
- `interval` support/truncation interval $[a, b]$
- `D` the minimum distance at degree `m`
- `convergence` An integer code. 0 indicates successful completion(the EM iteration is convergent). 1 indicates that the iteration limit `maxit` had been reached in the EM iteration;

made.mvar

Minimum Approximate Distance Estimate of Multivariate Density Function

Description

Minimum Approximate Distance Estimate of Multivariate Density Function

Usage

```
made.mvar(
  x,
  M0 = 1L,
  M,
  search = TRUE,
  interval = NULL,
  mar.deg = TRUE,
  method = c("cd", "quadprog"),
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

<code>x</code>	an $n \times d$ matrix or data.frame of multivariate sample of size n
<code>M0</code>	a positive integer or a vector of d positive integers specify starting candidate degrees for searching optimal degrees.
<code>M</code>	a positive integer or a vector of d positive integers specify the maximum candidate or the given model degrees for the joint density.
<code>search</code>	logical, whether to search optimal degrees between <code>M0</code> and <code>M</code> or not but use <code>M</code> as the given model degrees for the joint density.
<code>interval</code>	a vector of two endpoints or a $2 \times d$ matrix, each column containing the endpoints of support/truncation interval for each marginal density. If missing, the i -th column is assigned as $c(\min(x[, i]), \max(x[, i]))$.
<code>mar.deg</code>	logical, if TRUE, the optimal degrees are selected based on marginal data, otherwise, the optimal degrees are chosen the joint data. See details.
<code>method</code>	method for finding minimum distance estimate. "cd": coordinate-descent;
<code>controls</code>	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion <code>eps</code> . Default is <code>mable.ctrl</code> . See Details.
<code>progress</code>	if TRUE a text progressbar is displayed

Details

A d -variate density f on a hyperrectangle $[a, b] = [a_1, b_1] \times \cdots \times [a_d, b_d]$ can be approximated by a mixture of d -variate beta densities on $[a, b]$, $\beta_{m_j}(x) = \prod_{i=1}^d \beta_{m_i, j_i}[(x_i - a_i)/(b_i - a_i)]/(b_i - a_i)$, with proportion $p(j_1, \dots, j_d)$, $0 \leq j_i \leq m_i, i = 1, \dots, d$. If `search=TRUE` then the model degrees are chosen using a method of change-point based on the marginal data if `mar.deg=TRUE` or the joint data if `mar.deg=FALSE`. If `search=FALSE`, then the model degree is specified by `M`. For large data and multimodal density, the search for the model degrees is very time-consuming. In this case, it is suggested that use `method="cd"` and select the degrees based on marginal data using `mable` or `optimable`.

Value

A list with components

- `m` a vector of the selected optimal degrees by the method of change-point
- `p` a vector of the mixture proportions $p(j_1, \dots, j_d)$, arranged in the column-major order of $j = (j_1, \dots, j_d)$, $0 \leq j_i \leq m_i, i = 1, \dots, d$.
- `minD` the minimum distance at an optimal degree `m`
- `pval` the p-values of change-points for choosing the optimal degrees for the marginal densities
- `M` the vector (m_1, m_2, \dots, m_d) , where m_i is the largest candidate degree when the search stopped for the i -th marginal density
- `interval` support hyperrectangle $[a, b] = [a_1, b_1] \times \cdots \times [a_d, b_d]$
- `convergence` An integer code. 0 indicates successful completion(the EM iteration is convergent). 1 indicates that the iteration limit `maxit` had been reached in the EM iteration;

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. (2016) Efficient and robust density estimation using Bernstein type polynomials. *Journal of Nonparametric Statistics*, 28(2):250-271.

Wang, T. and Guan, Z.,(2019) Bernstein Polynomial Model for Nonparametric Multivariate Density, *Statistics*, Vol. 53, no. 2, 321-338

See Also

[mable](#), [optimable](#)

Examples

```
## Old Faithful Data

library(mable)
a<-c(0, 40); b<-c(7, 110)
ans<- made.mvar(faithful, M = c(46,19), search =FALSE, method="quadprog",
interval = rbind(a,b), progress=FALSE)
plot(ans, which="density")
plot(ans, which="cumulative")
```

madem.copula

Minimum Approximate Distance Estimate of Copula with given model degrees

Description

Minimum Approximate Distance Estimate of Copula with given model degrees

Usage

```
madem.copula(u, m)
```

Arguments

u an $n \times d$ matrix of (pseudo) observations.
m d-vector of model degrees

Details

With given model degrees m , the parameters p , the mixing proportions of the beta distribution, are calculated as the minimizer of the approximate L_2 distance between the empirical distribution and the Bernstein polynomial model.

Value

An invisible mable object with components

- m the given degree
- p the estimated vector of mixture proportions $p = (p_0, \dots, p_m)$ with the given degree m
- D the minimum distance at degree m

madem.density	<i>Minimum Approximate Distance Estimate of univariate Density Function with given model degree(s)</i>
---------------	--

Description

Minimum Approximate Distance Estimate of univariate Density Function with given model degree(s)

Usage

```
madem.density(
  x,
  m,
  p = rep(1, prod(m + 1))/prod(m + 1),
  interval = NULL,
  method = c("qp", "em"),
  maxit = 10000,
  eps = 1e-07
)
```

Arguments

x	an n x d matrix or data.frame of multivariate sample of size n
m	a positive integer or a vector of d positive integers specify the given model degrees for the joint density.
p	initial guess of p
interval	a vector of two endpoints or a 2 x d matrix, each column containing the endpoints of support/truncation interval for each marginal density. If missing, the i-th column is assigned as $c(\min(x[, i]), \max(x[, i]))$.
method	method for finding minimum distance estimate. "em": EM like method;
maxit	the maximum iterations
eps	the criterion for convergence

Details

A d -variate cdf F on a hyperrectangle $[a, b] = [a_1, b_1] \times \cdots \times [a_d, b_d]$ can be approximated by a mixture of d -variate beta cdfs on $[a, b]$, $\beta_{m,j}(x) = \prod_{i=1}^d B_{m_i, j_i}[(x_i - a_i)/(b_i - a_i)]$, with proportion $p(j_1, \dots, j_d)$, $0 \leq j_i \leq m_i, i = 1, \dots, d$. With a given model degree m , the parameters p , the mixing proportions of the beta distribution, are calculated as the minimizer of the approximate L_2 distance between the empirical distribution and the Bernstein polynomial model. The quadratic programming with linear constraints is used to solve the problem.

Value

An invisible mable object with components

- m the given model degree(s)
- p the estimated vector of mixture proportions with the given optimal degree(s) m
- interval support/truncation interval $[a, b]$
- D the minimum distance at degree m

maple.aft

Mable fit of AFT model with given regression coefficients

Description

Maximum approximate profile likelihood estimation of Bernstein polynomial model in accelerated failure time based on interval censored event time data with given regression coefficients which are efficient estimates provided by other semiparametric methods.

Usage

```
maple.aft(
  formula,
  data,
  M,
  g,
  tau = NULL,
  p = NULL,
  x0 = NULL,
  controls = maple.ctrl(),
  progress = TRUE
)
```

Arguments

formula	regression formula. Response must be cbind. See 'Details'.
data	a data frame containing variables in formula.

M	a positive integer or a vector (m0, m1). If M = m0 or m0 = m1, then m0 is a pre-selected degree. If m0 < m1 it specifies the set of consecutive candidate model degrees m0:m1 for searching an optimal degree, where m1-m0 > 3.
g	the given d -vector of regression coefficients.
tau	the right endpoint of the support or truncation interval $[0, \tau)$ of the baseline density. Default is NULL (unknown), otherwise if tau is given then it is taken as a known value of τ . See 'Details'.
p	an initial coefficients of Bernstein polynomial of degree m0, default is the uniform initial.
x0	a data frame specifying working baseline covariates on the right-hand-side of formula. See 'Details'.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

Consider the accelerated failure time model with covariate for interval-censored failure time data: $S(t|x) = S(t \exp(\gamma^T(x - x_0))|x_0)$, where x and x_0 may contain dummy variables and interaction terms. The working baseline x_0 in arguments contains only the values of terms excluding dummy variables and interaction terms in the right-hand-side of formula. Thus g is the initial guess of the coefficients γ of $x - x_0$ and could be longer than x_0 . Let $f(t|x)$ and $F(t|x) = 1 - S(t|x)$ be the density and cumulative distribution functions of the event time given $X = x$, respectively. Then $f(t|x_0)$ on a support or truncation interval $[0, \tau]$ can be approximated by $f_m(t|x_0; p) = \tau^{-1} \sum_{i=0}^m p_i \beta_{mi}(t/\tau)$, where $p_i \geq 0$, $i = 0, \dots, m$, $\sum_{i=0}^m p_i = 1$, $\beta_{mi}(u)$ is the beta density with shapes $i + 1$ and $m - i + 1$, and τ is larger than the largest observed time, either uncensored time, or right endpoint of interval/left censored, or left endpoint of right censored time. We can approximate $S(t|x_0)$ on $[0, \tau]$ by $S_m(t|x_0; p) = \sum_{i=0}^m p_i \bar{B}_{mi}(t/\tau)$, where $\bar{B}_{mi}(u)$ is the beta survival function with shapes $i + 1$ and $m - i + 1$.

Response variable should be of the form `cbind(1, u)`, where $(1, u)$ is the interval containing the event time. Data is uncensored if $1 = u$, right censored if $u = \text{Inf}$ or $u = \text{NA}$, and left censored data if $1 = 0$. The truncation time τ and the baseline x_0 should be chosen so that $S(t|x) = S(t \exp(\gamma^T(x - x_0))|x_0)$ on $[\tau, \infty)$ is negligible for all the observed x .

The search for optimal degree m stops if either m_1 is reached or the test for change-point results in a p-value `pval` smaller than `sig.level`.

Value

A list with components

- `m` the selected optimal degree m
- `p` the estimate of $p = (p_0, \dots, p_m)$, the coefficients of Bernstein polynomial of degree m
- `coefficients` the given regression coefficients of the AFT model
- `SE` the standard errors of the estimated regression coefficients
- `z` the z-scores of the estimated regression coefficients

- `mloglik` the maximum log-likelihood at an optimal degree `m`
- `tau.n` maximum observed time τ_n
- `tau` right endpoint of truncation interval $[0, \tau)$
- `x0` the working baseline covariates
- `egx0` the value of $e^{\gamma^T x_0}$
- `convergence` an integer code, 1 indicates either the EM-like iteration for finding maximum likelihood reached the maximum iteration for at least one `m` or the search of an optimal degree using change-point method reached the maximum candidate degree, 2 indicates both occurred, and 0 indicates a successful completion.
- `delta` the final `delta` if `m0 = m1` or the final `pval` of the change-point for searching the optimal degree `m`;

and, if `m0 < m1`,

- `M` the vector (m_0, m_1) , where `m1` is the last candidate when the search stopped
- `lk` log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- `lr` likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- `pval` the p-values of the change-point tests for choosing optimal model degree
- `chpts` the change-points chosen with the given candidate model degrees

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. (2019) Maximum Approximate Likelihood Estimation in Accelerated Failure Time Model for Interval-Censored Data, arXiv:1911.07087.

See Also

[mable.aft](#)

Examples

```
## Breast Cosmesis Data
g<-0.41 #Hanson and Johnson 2004, JCGS,
res1<-maple.aft(cbind(left, right)~treat, data=cosmesis, M=c(1,30), g=g,
               tau=100, x0=data.frame(treat="RCT"))
op<-par(mfrow=c(1,2), lwd=1.5)
plot(x=res1, which="likelihood")
plot(x=res1, y=data.frame(treat="RT"), which="survival", model='aft', type="l", col=1,
     add=FALSE, main="Survival Function")
plot(x=res1, y=data.frame(treat="RCT"), which="survival", model='aft', lty=2, col=1)
legend("bottomleft", bty="n", lty=1:2, col=1, c("Radiation Only", "Radiation and Chemotherapy"))
par(op)
```

maple.dr	<i>Maximum approximate profile likelihood estimate of the density ratio model</i>
----------	---

Description

Select optimal degree with a given regression coefficients.

Usage

```
maple.dr(
  x,
  y,
  M,
  regr,
  ...,
  interval = c(0, 1),
  alpha = NULL,
  vb = 0,
  baseline = NULL,
  controls = mable.ctrl(),
  progress = TRUE,
  message = TRUE
)
```

Arguments

x, y	original two sample raw data, x:"Control", y: "Case".
M	a positive integer or a vector (m0, m1).
regr	regressor vector function $r(x) = (1, r_1(x), \dots, r_d(x))$ which returns n x (d+1) matrix, n=length(x)
...	additional arguments to be passed to regr
interval	a vector (a,b) containing the endpoints of supporting/truncation interval of x and y.
alpha	a given regression coefficient, missing value is imputed by logistic regression
vb	code for vanishing boundary constraints, -1: f0(a)=0 only, 1: f0(b)=0 only, 2: both, 0: none (default).
baseline	the working baseline, "Control" or "Case", if NULL it is chosen to the one with smaller estimated lower bound for model degree.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion for EM and Newton iterations. Default is <code>mable.ctrl</code> . See Details.
progress	logical: should a text progressbar be displayed
message	logical: should warning messages be displayed

Details

Suppose that ("control") and y ("case") are independent samples from f_0 and f_1 which satisfy $f_1(x) = f_0(x) \exp[\alpha_0 + \alpha' r(x)]$ with $r(x) = (r_1(x), \dots, r_d(x))$. Maximum approximate Bernstein/Beta likelihood estimates of f_0 and f_1 are calculated with a given regression coefficients which are efficient estimates provided by other semiparametric methods such as logistic regression. If support is (a, b) then replace $r(x)$ by $r[a + (b-a)x]$. For a fixed m , using the Bernstein polynomial model for baseline f_0 , MABLEs of f_0 and parameters α can be estimated by EM algorithm and Newton iteration. If estimated lower bound m_b for m based on y is smaller than that based on x , then switch x and y and f_1 is used as baseline. If $M=m$ or $m_0=m_1=m$, then m is a preselected degree. If $m_0 < m_1$ it specifies the set of consecutive candidate model degrees $m_0:m_1$ for searching an optimal degree by the change-point method, where $m_1 - m_0 > 3$.

Value

A list with components

- m the given or a selected degree by method of change-point
- p the estimated vector of mixture proportions $p = (p_0, \dots, p_m)$ with the given or selected degree m
- α the given regression coefficients
- $mloglik$ the maximum log-likelihood at degree m
- $interval$ support/truncation interval (a, b)
- $baseline = "control"$ if f_0 is used as baseline, or $"case"$ if f_1 is used as baseline.
- M the vector (m_0, m_1) , where m_1 , if greater than m_0 , is the largest candidate when the search stopped
- lk log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- lr likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- $pval$ the p-values of the change-point tests for choosing optimal model degree
- $chpts$ the change-points chosen with the given candidate model degrees

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z., Maximum Approximate Bernstein Likelihood Estimation of Densities in a Two-sample Semiparametric Model

maple.dr.group	<i>Maximum approximate profile likelihood estimate of the density ratio model for grouped data with given regression coefficients</i>
----------------	---

Description

Select optimal degree of Bernstein polynomial model for grouped data with a given regression coefficients.

Usage

```
maple.dr.group(
  t,
  n0,
  n1,
  M,
  regr,
  ...,
  interval = c(0, 1),
  alpha = NULL,
  vb = 0,
  controls = mable.ctrl(),
  progress = TRUE,
  message = TRUE
)
```

Arguments

t	cutpoints of class intervals
n0, n1	frequencies of two sample data grouped by the classes specified by t. n0:"Control", n1: "Case".
M	a positive integer or a vector (m0, m1).
regr	regressor vector function $r(x) = (1, r_1(x), \dots, r_d(x))$ which returns $n \times (d+1)$ matrix, $n=\text{length}(x)$
...	additional arguments to be passed to regr
interval	a vector (a,b) containing the endpoints of supporting/truncation interval of x and y.
alpha	a given regression coefficient, missing value is imputed by logistic regression
vb	code for vanishing boundary constraints, -1: $f_0(a)=0$ only, 1: $f_0(b)=0$ only, 2: both, 0: none (default).
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion for EM and Newton iterations. Default is <code>mable.ctrl</code> . See Details.
progress	logical: should a text progressbar be displayed
message	logical: should warning messages be displayed

Details

Suppose that n_0 ("control") and n_1 ("case") are frequencies of independent samples grouped by the classes t from f_0 and f_1 which satisfy $f_1(x)=f_0(x)\exp[\alpha_0+\alpha'r(x)]$ with $r(x)=(r_1(x),\dots,r_d(x))$. Maximum approximate Bernstein/Beta likelihood estimates of f_0 and f_1 are calculated with a given regression coefficients which are efficient estimates provided by other semiparametric methods such as logistic regression. If support is (a,b) then replace $r(x)$ by $r[a+(b-a)x]$. For a fixed m , using the Bernstein polynomial model for baseline f_0 , MABLEs of f_0 and parameters α can be estimated by EM algorithm and Newton iteration. If estimated lower bound m_b for m based on n_1 is smaller than that based on n_0 , then switch n_0 and n_1 and use f_1 as baseline. If $M=m$ or $m_0=m_1=m$, then m is a preselected degree. If $m_0 < m_1$ it specifies the set of consecutive candidate model degrees $m_0:m_1$ for searching an optimal degree by the change-point method, where $m_1 - m_0 > 3$.

Value

A list with components

- m the given or a selected degree by method of change-point
- p the estimated vector of mixture proportions $p = (p_0, \dots, p_m)$ with the given or selected degree m
- α the given regression coefficients
- $mloglik$ the maximum log-likelihood at degree m
- $interval$ support/truncation interval (a, b)
- $baseline$ ="control" if f_0 is used as baseline, or ="case" if f_1 is used as baseline.
- M the vector (m_0, m_1) , where m_1 , if greater than m_0 , is the largest candidate when the search stopped
- lk log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- lr likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- $pval$ the p-values of the change-point tests for choosing optimal model degree
- $chpts$ the change-points chosen with the given candidate model degrees

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z., Application of Bernstein Polynomial Model to Density and ROC Estimation in a Semi-parametric Density Ratio Model

maple.ph

Mable fit of the PH model with given regression coefficients

Description

Maximum approximate profile likelihood estimation of Bernstein polynomial model in Cox's proportional hazards regression based on interval censored event time data with given regression coefficients which are efficient estimates provided by other semiparametric methods.

Usage

```
maple.ph(
  formula,
  data,
  M,
  g,
  pi0 = NULL,
  p = NULL,
  tau = Inf,
  x0 = NULL,
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

formula	regression formula. Response must be cbind. See 'Details'.
data	a data frame containing variables in formula.
M	a positive integer or a vector (m0, m1). If M = m0 or m0 = m1, then m0 is a pre-selected degree. If m0 < m1 it specifies the set of consecutive candidate model degrees m0:m1 for searching an optimal degree, where m1-m0>3.
g	the given <i>d</i> -vector of regression coefficients
pi0	Initial guess of $\pi(x_0) = F(\tau_n x_0)$. Without right censored data, pi0 = 1. See 'Details'.
p	an initial coefficients of Bernstein polynomial model of degree m0, default is the uniform initial.
tau	right endpoint of support $[0, \tau)$ must be greater than or equal to the maximum observed time
x0	a data frame specifying working baseline covariates on the right-hand-side of formula. See 'Details'.
controls	Object of class mable.ctrl() specifying iteration limit and other control options. Default is mable.ctrl.
progress	if TRUE a text progressbar is displayed

Details

Consider Cox's PH model with covariate for interval-censored failure time data: $S(t|x) = S(t|x_0)\exp(\gamma^T(x-x_0))$, where x_0 satisfies $\gamma^T(x-x_0) \geq 0$, where x and x_0 may contain dummy variables and interaction terms. The working baseline x_0 in arguments contains only the values of terms excluding dummy variables and interaction terms in the right-hand-side of formula. Thus g is the initial guess of the coefficients γ of $x-x_0$ and could be longer than x_0 . Let $f(t|x)$ and $F(t|x) = 1 - S(t|x)$ be the density and cumulative distribution functions of the event time given $X = x$, respectively. Then $f(t|x_0)$ on $[0, \tau_n]$ can be approximated by $f_m(t|x_0; p) = \tau_n^{-1} \sum_{i=0}^m p_i \beta_{mi}(t/\tau_n)$, where $p_i \geq 0$, $i = 0, \dots, m$, $\sum_{i=0}^m p_i = 1 - p_{m+1}$, $\beta_{mi}(u)$ is the beta density with shapes $i+1$ and $m-i+1$, and τ_n is the largest observed time, either uncensored time, or right endpoint of interval/left censored, or left endpoint of right censored time. So we can approximate $S(t|x_0)$ on $[0, \tau_n]$ by $S_m(t|x_0; p) = \sum_{i=0}^{m+1} p_i \bar{B}_{mi}(t/\tau_n)$, where $\bar{B}_{mi}(u)$, $i = 0, \dots, m$, is the beta survival function with shapes $i+1$ and $m-i+1$, $\bar{B}_{m,m+1}(t) = 1$, $p_{m+1} = 1 - \pi(x_0)$, and $\pi(x_0) = F(\tau_n|x_0)$. For data without right-censored time, $p_{m+1} = 1 - \pi(x_0) = 0$.

Response variable should be of the form `cbind(1, u)`, where $(1, u)$ is the interval containing the event time. Data is uncensored if $1 = u$, right censored if $u = \text{Inf}$ or $u = \text{NA}$, and left censored data if $1 = \emptyset$. The associated covariate contains d columns. The baseline x_0 should be chosen so that $\gamma^T(x-x_0)$ is nonnegative for all the observed x .

The search for optimal degree m stops if either m_1 is reached or the test for change-point results in a p-value `pval` smaller than `sig.level`.

Value

a class 'mable_reg' object, a list with components

- `M` the vector (m_0, m_1) , where m_1 is the last candidate degree when the search stopped
- `m` the selected optimal degree m
- `p` the estimate of $p = (p_0, \dots, p_m, p_{m+1})$, the coefficients of Bernstein polynomial of degree m
- `coefficients` the given regression coefficients of the PH model
- `mloglik` the maximum log-likelihood at an optimal degree m
- `lk` log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- `lr` likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- `tau.n` maximum observed time τ_n
- `tau` right endpoint of support $[0, \tau)$
- `x0` the working baseline covariates
- `egx0` the value of $e^{\gamma^T x_0}$
- `convergence` an integer code. 0 indicates successful completion (the iteration is convergent). 1 indicates that the maximum candidate degree had been reached in the calculation;
- `delta` the final convergence criterion for EM iteration;
- `chpts` the change-points among the candidate degrees;
- `pom` the p-value of the selected optimal degree m as a change-point;

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. (2019) Maximum Approximate Bernstein Likelihood Estimation in Proportional Hazard Model for Interval-Censored Data, arXiv:1906.08882 .

See Also

[mable.ph](#)

Examples

```
## Simulated Weibull data
require(icenReg)
set.seed(123)
simdata<-simIC_weib(70, inspections = 5, inspectLength = 1)
sp<-ic_sp(cbind(1, u) ~ x1 + x2, data = simdata)
res0<-mable.ph(cbind(1, u) ~ x1 + x2, data = simdata, M=c(2,20),
              g=sp$coefficients, tau=7)
op<-par(mfrow=c(1,2))
plot(res0, which=c("likelihood","change-point"))
par(op)
res1<-mable.ph(cbind(1, u) ~ x1 + x2, data = simdata, M=res0$m,
              g=c(.5,-.5), tau=7)
op<-par(mfrow=c(1,2))
plot(res1, y=data.frame(x1=0, x2=0), which="density", add=FALSE, type="l",
      xlab="Time", main="Desnity Function")
lines(xx<-seq(0, 7, len=512), dweibull(xx, 2,2), lty=2, col=2)
legend("topright", bty="n", lty=1:2, col=1:2, c("Estimated","True"))
plot(res1, y=data.frame(x1=0, x2=0), which="survival", add=FALSE, type="l",
      xlab="Time", main="Survival Function")
lines(xx, 1-pweibull(xx, 2, 2), lty=2, col=2)
legend("topright", bty="n", lty=1:2, col=1:2, c("Estimated","True"))
par(op)
```

maple.po

Mable fit of the PO model with given regression coefficients

Description

Maximum approximate profile likelihood estimation of Bernstein polynomial model in proportional odds rate regression based on interal censored event time data with given regression coefficients and select an optimal degree m if coefficients are efficient estimates provided by other semiparametric methods.

Usage

```
maple.po(
  formula,
  data,
  M,
  g,
  tau,
  x0 = NULL,
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

formula	regression formula. Response must be cbind. See 'Details'.
data	a data frame containing variables in formula.
M	a positive integer or a vector (m0, m1). If M = m or m0 = m1 = m, then m is a pre-selected degree. If m0 < m1 it specifies the set of consecutive candidate model degrees m0:m1 for searching an optimal degree, where m1 - m0 > 3.
g	the given <i>d</i> -vector of regression coefficients
tau	right endpoint of support $[0, \tau]$ must be greater than or equal to the maximum observed time
x0	a data frame specifying working baseline covariates on the right-hand-side of formula. See 'Details'.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

Consider Generalized PO model with covariate for interval-censored failure time data: $S(t|x) = S(t|x_0)^{\exp(\gamma'(x-x_0))}$, where x_0 satisfies $\gamma'(x - x_0) \geq 0$, where x and x_0 may contain dummy variables and interaction terms. The working baseline `x0` in arguments contains only the values of terms excluding dummy variables and interaction terms in the right-hand-side of `formula`. Thus `g` is the initial guess of the coefficients γ of $x - x_0$ and could be longer than `x0`. Let $f(t|x)$ and $F(t|x) = 1 - S(t|x)$ be the density and cumulative distribution functions of the event time given $X = x$, respectively. Then $f(t|x_0)$ on $[0, \tau_n]$ can be approximated by $f_m(t|x_0; p) = \tau^{-1} \sum_{i=0}^m p_i \beta_{mi}(t/\tau)$, where $p_i \geq 0$, $i = 0, \dots, m$, $\sum_{i=0}^m p_i = 1$, $\beta_{mi}(u)$ is the beta density with shapes $i + 1$ and $m - i + 1$, and τ is the right endpoint of support interval. So we can approximate $S(t|x_0)$ on $[0, \tau]$ by $S_m(t|x_0; p) = \sum_{i=0}^m p_i \bar{B}_{mi}(t/\tau)$, where $\bar{B}_{mi}(u)$, $i = 0, \dots, m$, is the beta survival function with shapes $i + 1$ and $m - i + 1$.

Response variable should be of the form `cbind(1, u)`, where $(1, u)$ is the interval containing the event time. Data are uncensored if $1 = u$, right censored if $u = \text{Inf}$ or $u = \text{NA}$, and left censored data if $1 = 0$. The associated covariate contains *d* columns. The baseline `x0` should be chosen so that $\gamma'(x - x_0)$ is nonnegative for all the observed x .

The search for optimal degree *m* stops if either *m1* is reached or the test for change-point results in a p-value `pval` smaller than `sig.level`.

Value

a class 'mable_reg' object, a list with components

- M the vector (m_0, m_1) , where m_1 is the last candidate degree when the search stopped
- m the selected optimal degree m
- p the estimate of $p = (p_0, \dots, p_m, p_{m+1})$, the coefficients of Bernstein polynomial of degree m
- coefficients the given regression coefficients of the PH model
- mloglik the maximum log-likelihood at an optimal degree m
- lk log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- lr likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- tau.n maximum observed time τ_n
- tau right endpoint of support $[0, \tau)$
- x0 the working baseline covariates
- egx0 the value of $e^{\gamma'x_0}$
- convergence an integer code, 0 indicates successful completion(the iteration is convergent), 1 indicates that the maximum candidate degree had been reached in the calculation;
- delta the final convergence criterion for EM iteration;
- chpts the change-points among the candidate degrees;
- pom the p-value of the selected optimal degree m as a change-point;

Author(s)

Zhong Guan <zguan@iu.edu>

References

Guan, Z. et al. (???) Maximum Approximate Bernstein Likelihood Estimation in Generalized Proportional Odds Regression Model for Interval-Censored Data

See Also

[mable.po](#)

Examples

```
## Simulated Weibull data
require(icenReg)
set.seed(111)
simdata<-simIC_weib(100, model = "po", inspections = 2,
  inspectLength = 2.5, prob_cen=1)
sp<-ic_sp(cbind(1, u) ~ x1 + x2, data = simdata, model="po")
gt<--sp$coefficients
res0<-maple.po(cbind(1, u) ~ x1 + x2, data = simdata, M=c(1,20), g=gt, tau=6)
op<-par(mfrow=c(1,2))
plot(res0, which=c("likelihood", "change-point"))
```

```

par(op)
res1<-mable.po(cbind(1, u) ~ x1 + x2, data = simdata, M=c(1,20),
  g=gt, tau=6, x0=data.frame(x1=max(simdata$x1),x2=-1))
op<-par(mfrow=c(2,2))
plot(res1, which=c("likelihood","change-point"))
plot(res0, y=data.frame(x1=0,x2=0), which="density", add=FALSE, type="l",
  xlab="Time", main="Desnity Function")
plot(res1, y=data.frame(x1=0,x2=0), which="density", add=TRUE, lty=2, col=4)
lines(xx<-seq(0, 7, len=512), dweibull(xx, 2,2), lty=3, col=2, lwd=1.5)
legend("topright", bty="n", lty=1:3, col=c(1,4,2), c(expression(hat(f)[0]),
  expression(tilde(f)[0]), expression(f[0])))
plot(res0, y=data.frame(x1=0,x2=0), which="survival", add=FALSE, type="l",
  xlab="Time", main="Survival Function")
plot(res1, y=data.frame(x1=0,x2=0), which="survival", add=TRUE, lty=2, col=4)
lines(xx, 1-pweibull(xx, 2, 2), lty=2, col=2)
legend("topright", bty="n", lty=1:3, col=c(1,4,2), c(expression(hat(S)[0]),
  expression(tilde(S)[0]), expression(S[0])))
par(op)

```

marginal.p

The mixing proportions of marginal distribution from the mixture of multivariate beta distribution

Description

The mixing proportions of marginal distribution from the mixture of multivariate beta distribution

Usage

```
marginal.p(p, m)
```

Arguments

p	the mixing proportions of the mixture of multivariate beta distribution
m	the model degrees $m=(m_1, \dots, m_d)$ of the mixture of multivariate beta distribution

Value

a list of mixing proportions of all the marginal distributions

mvecdf	<i>Multivariate empirical cumulative distribution evaluated at sample data</i>
--------	--

Description

Multivariate empirical cumulative distribution evaluated at sample data

Usage

mvecdf(x)

Arguments

x an n x d matrix of data values, rows are n observations of $X = (X_1, \dots, X_d)$

Value

a vector of n values

mvpbeta	<i>Component Beta cumulative distribution functions of the Bernstein polynomial model</i>
---------	---

Description

Component Beta cumulative distribution functions of the Bernstein polynomial model

Usage

mvpbeta(x, m)

Arguments

x n x d matrix, rows are n observations of $X=(X_1,\dots,X_d)$

m vector of d nonnegative integers $m=(m[1], \dots, m[d])$.

Value

an n x K matrix, $K=(m[1]+1) \dots (m[d]+1)$.

 optim.gcp

Choosing optimal model degree by gamma change-point method

Description

Choose an optimal degree using gamma change-point model with two changing shape and scale parameters.

Usage

```
optim.gcp(obj)
```

Arguments

obj a class "mable" or 'mable_reg' object containing a vector $M = (m_0, m_1)$, lk, log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$

Value

a list with components

- m the selected optimal degree m
- M the vector (m_0, m_1) , where m_1 is the last candidate when the search stopped
- mloglik the maximum log-likelihood at degree m
- interval support/truncation interval (a, b)
- lk log-likelihoods evaluated at $m \in \{m_0, \dots, m_1\}$
- lr likelihood ratios for change-points evaluated at $m \in \{m_0 + 1, \dots, m_1\}$
- pval the p-values of the change-point tests for choosing optimal model degree
- chpts the change-points chosen with the given candidate model degrees

Examples

```
# simulated data
p<-c(1:5,5:1)
p<-p/sum(p)
x<-rmixbeta(100, p)
res1<-mable(x, M=c(2, 50), IC="none")
m1<-res1$m[1]
res2<-optim.gcp(res1)
m2<-res2$m
op<-par(mfrow=c(1,2))
plot(res1, which="likelihood", add=FALSE)
plot(res2, which="likelihood")
#segments(m2, min(res1$lk), m2, res2$mloglik, col=4)
plot(res1, which="change-point", add=FALSE)
plot(res2, which="change-point")
par(op)
```

optimable	<i>mable with degree selected by the method of moment and method of mode</i>
-----------	--

Description

Maximum Approximate Bernstein/Beta Likelihood Estimation with an optimal model degree estimated by the Method of Moment

Usage

```
optimable(
  x,
  interval,
  m = NULL,
  mu = NULL,
  lam = NULL,
  modes = NULL,
  nmod = 1,
  ushaped = FALSE,
  maxit = 50L
)
```

Arguments

x	a univariate sample data in interval
interval	a closed interval $c(a, b)$, default is $[0, 1]$
m	initial degree, default is 2 times the number of modes nmod.
mu	a vector of component means of multimodal mixture density, default is NULL for unimodal or unknown
lam	a vector of mixture proportions of same length of mu
modes	a vector of the locations of modes, if it is NULL (default) and <code>multimode::locmodes()</code>
nmod	the number of modes, if nmod=0, the lower bound for m is estimated based on mean and variance only.
ushaped	logical, whether or not the density is clearly U-shaped including J- and L-shaped with mode occurs at the endpoint of the support.
maxit	maximum iterations

Details

If the data show a clear uni- or multi-modal distribution, then give the value of nmod as the number of modes. Otherwise nmod=0. The degree is estimated by the iterative method of moment with an initial degree estimated by the method of mode. For multimodal density, if useful estimates of the component means mu and proportions lam are available then they can be used to give an initial degree. If the distribution is clearly U-, J-, or L-shaped, i.e., the mode occurs at the endpoint of interval, then set ushaped=TRUE. In this case the degree is estimated by the method of mode.

Value

A class "mable" object with components

- `m` the given or a selected degree by method of change-point
- `p` the estimated vector of mixture proportions $p = (p_0, \dots, p_m)$ with the selected/given optimal degree `m`
- `mloglik` the maximum log-likelihood at degree `m`
- `interval` support/truncation interval `(a, b)`
- `convergence` An integer code. 0 indicates successful completion (all the EM iterations are convergent and an optimal degree is successfully selected in `M`). Possible error codes are
 - 1, indicates that the iteration limit `maxit` had been reached in at least one EM iteration;
 - 2, the search did not finish before `m1`.
- `delta` the convergence criterion `delta` value

Author(s)

Zhong Guan <zguan@iu.edu>

Examples

```
## Old Faithful Data
x<-faithful
x1<-faithful[,1]
x2<-faithful[,2]
a<-c(0, 40); b<-c(7, 110)
mu<-(apply(x,2,mean)-a)/(b-a)
s2<-apply(x,2,var)/(b-a)^2
# mixing proportions
lambda<-c(mean(x1<3),mean(x2<65))
# guess component mean
mu1<-(c(mean(x1[x1<3]), mean(x2[x2<65]))-a)/(b-a)
mu2<-(c(mean(x1[x1>=3]), mean(x2[x2>=65]))-a)/(b-a)
# estimate lower bound for m
mb<-ceiling((mu*(1-mu)-s2)/(s2-lambda*(1-lambda)*(mu1-mu2)^2)-2)
mb
m1<-optimable(x1, interval=c(a[1],b[1]), nmod=2, modes=c(2,4.5))$m
m2<-optimable(x2, interval=c(a[2],b[2]), nmod=2, modes=c(52.5,80))$m
m1;m2
erupt1<-mable(x1, M=mb[1], interval=c(a[1],b[1]))
erupt2<-mable(x1, M=m1, interval=c(a[1],b[1]))
wait1<-mable(x2, M=mb[2],interval=c(a[2],b[2]))
wait2<-mable(x2, M=m2,interval=c(a[2],b[2]))
ans1<- mable.mvar(faithful, M = mb, search =FALSE, method="em", interval = cbind(a,b))
ans2<- mable.mvar(faithful, M = c(m1,m2), search =FALSE, method="em", interval = cbind(a,b))
op<-par(mfrow=c(1,2), cex=0.8)
hist(x1, probability = TRUE, col="grey", border="white", main="",
      xlab="Eruptions", ylim=c(0,.65), las=1)
plot(erupt1, add=TRUE,"density")
plot(erupt2, add=TRUE,"density",lty=2,col=2)
```

```

legend("topleft", lty=c(1,2),col=1:2, bty="n", cex=.7,
      c(expression(paste("m = ", m[b])),expression(paste("m = ", hat(m)))))
hist(x2, probability = TRUE, col="grey", border="white", main="",
     xlab="Waiting", las=1)
plot(wait1, add=TRUE,"density")
plot(wait2, add=TRUE,"density",lty=2,col=2)
legend("topleft", lty=c(1,2),col=1:2, bty="n", cex=.7,
      c(expression(paste("m = ", m[b])),expression(paste("m = ", hat(m)))))
par(op)
op<-par(mfrow=c(1,2), cex=0.7)
plot(ans1, which="density", contour=TRUE)
plot(ans2, which="density", contour=TRUE, add=TRUE, lty=2, col=2)
plot(ans1, which="cumulative", contour=TRUE)
plot(ans2, which="cumulative", contour=TRUE, add=TRUE, lty=2, col=2)
par(op)

```

pancreas

Pancreatic Cancer Biomarker Data

Description

Contain sera measurements from 51 control patients with pancreatitis and 90 case patients with pancreatic cancer at the Mayo Clinic with a cancer antigen, CA125, and with a carbohydrate antigen, CA19-9 (Wieand, et al, 1989)

Usage

```
data(pancreas)
```

Format

A data frame with 141 rows and 3 variables.

- ca199. CA19-9 levels
- ca125. CA125 levels
- status. 0 = controls (non-cancer) and 1 = cases (cancer).

Source

Wieand, S., Gail, M. H., James, B. R., and James, K.L. (1989). A family of nonparametric statistics for comparing diagnostic markers with paired or unpaired data. *Biometrika*, 76, 585–592.

References

Wieand, S., Gail, M. H., James, B. R., and James, K.L. (1989). A family of nonparametric statistics for comparing diagnostic markers with paired or unpaired data. *Biometrika*, 76, 585–592.

Examples

```
data(pancreas)
```

plot.mable	<i>Plot method for class 'mable'</i>
------------	--------------------------------------

Description

Plot method for class 'mable'

Usage

```
## S3 method for class 'mable'
plot(
  x,
  which = c("density", "cumulative", "survival", "likelihood", "change-point", "all"),
  add = FALSE,
  contour = FALSE,
  lgd.x = NULL,
  lgd.y = NULL,
  nx = 512,
  ...
)
```

Arguments

x	Class "mable" object return by mablem, mable, mable.mvar, mablem.group or mable.group functions which contains p, mloglik, and M = m0:m1, lk, lr,
which	indicates which graphs to plot, options are "density", "cumulative", "likelihood", "change-point", "all". If not "all", which can contain more than one options.
add	logical add to an existing plot or not
contour	logical plot contour or not for two-dimensional data
lgd.x, lgd.y	coordinates of position where the legend is displayed
nx	number of evaluations of density, or cumulative distribution curve to be plotted.
...	additional arguments to be passed to the base plot function

Value

The data used for 'plot()', 'lines()', or 'persp()' are returned invisibly.

plot.mable_reg *Plot method for class 'mable_reg'*

Description

Plot method for class 'mable_reg'

Usage

```
## S3 method for class 'mable_reg'
plot(
  x,
  y,
  newdata = NULL,
  ntime = 512,
  xlab = "Time",
  which = c("survival", "likelihood", "change-point", "density", "all"),
  add = FALSE,
  ...
)
```

Arguments

x	a class 'mable_reg' object return by functions such as mable.ph which contains M, coefficients, p, m, x0, tau.n, tau.lk, lr.
y	a new data.frame of covariate value(s) as row(s), whose columns are arranged in the same order as in the formula called by the function that returned the object x.
newdata	a new data.frame (ignored if y is included), imputed by the working baseline x0 if both missing.
ntime	number of evaluations of density, survival or cumulative distribution curve to be plotted.
xlab	x-axis label
which	indicates which graphs to plot, options are "survival", "likelihood", "change-point", "density", or "all". If not "all", which can contain more than one options.
add	logical add to an existing plot or not
...	additional arguments to be passed to the base plot function

Author(s)

Zhong Guan <zguan@iu.edu>

se.coef.dr

Standard errors of coefficients in density ratio model

Description

Bootstrap estimates of standard errors for the regression coefficients which are estimated by maximum approximate Bernstein/Beta likelihood estimation method in a density ratio model based on two-sample raw data.

Usage

```
se.coef.dr(
  obj,
  grouped = FALSE,
  B = 500L,
  parallel = FALSE,
  ncore = NULL,
  controls = mable.ctrl()
)
```

Arguments

obj	Class 'mable_dr' object return by <code>mable.dr</code> or <code>mable.dr.group</code> functions
grouped	logical: are data grouped or not.
B	number of bootstrap runs.
parallel	logical: do parallel or not.
ncore	number of cores used for parallel computing. Default is half of availables.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and the convergence criterion for EM and Newton iterations. Default is <code>mable.ctrl</code> . See Details.

Details

Bootstrap method is used based on bootstrap samples generated from the MABLE's of the densities f_0 and f_1 . The bootstrap samples are fitted by the Bernstein polynomial model and the `glm()` to obtain bootstrap versions of coefficient estimates.

Value

the estimated standard errors

summary.mable *Summary methods for classes 'mable' and 'mable_reg'*

Description

Produces a summary of a mable fit.

Usage

```
## S3 method for class 'mable'
summary(object, ...)

## S3 method for class 'mable_reg'
summary(object, ...)
```

Arguments

object Class "mable" or 'mable_reg' object return by mable or mable.xxxx functions
 ... for future methods

Value

Invisibly returns its argument, object.

Examples

```
## Breast Cosmesis Data
aft.res<-mable.aft(cbind(left, right)~treat, data=cosmesis, M=c(1, 30), g=.41,
  tau=100, x0=data.frame(treat="RCT"))
summary(aft.res)
```

umc.mat *Matrix of the uniform marginal constraints*

Description

Matrix of the uniform marginal constraints

Usage

```
umc.mat(m)
```

Arguments

m vector of d nonnegative integers $m=(m[1], \dots, m[d])$.

Details

the matrix of the uniform marginal constraints A is used to form the linear equality constraints on parameter p : $Ap = 1/(m + 1)$.

Value

an $|m| \times K$ matrix, $|m|=m[1]+\dots+m[d]$, $K=(m[1]+1)\dots(m[d]+1)$.

 Vaal.Flow

Vaal River Annual Flow Data

Description

The annual flow data of Vaal River at Standerton as given by Table 1.1 of Linhart and Zucchini (1986) give the flow in millions of cubic metres.

Usage

```
data(Vaal.Flow)
```

Format

The format is: int [1:65] 222 1094 452 1298 882 988 276 216 103 490 ...

References

Linhart, H., and Zucchini, W., *Model Selection*, Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics, New York: John Wiley and Sons Inc, 1986.

Examples

```
data(Vaal.Flow)
```

 weib.gpo

Generalized PO model with Weibull baseline

Description

Maximum likelihood estimation in generalized proportional odds rate regression model with Weibull baseline based on interval censored event time data

Usage

```
weib.gpo(
  formula,
  data,
  g,
  scale,
  shape,
  eta = 1,
  eta.known = TRUE,
  controls = mable.ctrl(),
  progress = TRUE
)
```

Arguments

formula	regression formula. Response must be cbind. See 'Details'.
data	a dataset
g	initial d -vector of regression coefficients
scale	initial guess of the scale parameter for Weibull baseline
shape	initial guess of the shape parameter for Weibull baseline
eta	the given positive value of η . See 'Details'.
eta.known	logical. If TRUE eta is the known values of η , else eta is an initial guess of η . See 'Details'.
controls	Object of class <code>mable.ctrl()</code> specifying iteration limit and other control options. Default is <code>mable.ctrl</code> .
progress	if TRUE a text progressbar is displayed

Details

???

Value

a class 'mable_reg' object, a list with components

- convergence an integer code, 0 indicates successful completion(the iteration is convergent), 1 indicates that the maximum iteration had been reached in the calculation;
- delta the final convergence criterion for Newton iteration;

Examples

```
## Simulated Weibull data
require(icenReg)
set.seed(111)
simdata<-simIC_weib(100, model = "po", inspections = 2,
  inspectLength = 2.5, prob_cen=1)
sp<-ic_sp(cbind(1, u) ~ x1 + x2, data = simdata, model="po")
```



```

gt<--sp$coefficients
res0<-maple.po(cbind(1, u) ~ x1 + x2, data = simdata, M=c(1,20), g=gt, tau=6)
op<-par(mfrow=c(1,2))
plot(res0, which=c("likelihood","change-point"))
par(op)
res1<-mable.po(cbind(1, u) ~ x1 + x2, data = simdata, M=c(1,20), g=gt,
  tau=6, x0=data.frame(x1=max(simdata$x1),x2=-1))
res2<-weib.gpo(cbind(1, u) ~ x1 + x2, data = simdata, g=gt, scale=2, shape=2)
op<-par(mfrow=c(2,2))
plot(res1, which=c("likelihood","change-point"))
plot(res0, y=data.frame(x1=0,x2=0), which="density", add=FALSE, type="l",
  xlab="Time", main="Desnity Function")
plot(res1, y=data.frame(x1=0,x2=0), which="density", add=TRUE, lty=2, col=4)
lines(xx<-seq(0, 7, len=512), dweibull(xx, 2,2), lty=3, col=2, lwd=1.5)
lines(xx, dweibull(xx, res2$shape, res2$scale), lty=5, col=5, lwd=1.5)
legend("topright", bty="n", lty=1:3, col=c(1,4,2), c(expression(hat(f)[0]),
  expression(tilde(f)[0]), expression(f[0])))
plot(res0, y=data.frame(x1=0,x2=0), which="survival", add=FALSE, type="l",
  xlab="Time", main="Survival Function")
plot(res1, y=data.frame(x1=0,x2=0), which="survival", add=TRUE, lty=2, col=4)
lines(xx, 1-pweibull(xx, 2, 2), lty=2, col=2)
lines(xx, 1-pweibull(xx, res2$shape, res2$scale), lty=5, col=5, lwd=1.5)
legend("topright", bty="n", lty=1:3, col=c(1,4,2), c(expression(hat(S)[0]),
  expression(tilde(S)[0]), expression(S[0])))
par(op)

```

Index

- * **Accelerated failure time model**
 - mable.aft, 15
 - maple.aft, 49
- * **Additive measurement error**
 - mable.decon, 21
- * **Bernstein polynomial model**
 - dmixbeta, 7
 - dtmixbeta, 10
 - mable, 11
 - mable.decon, 21
 - mable.group, 27
- * **Cox proportional hazards model**
 - mable.ph, 36
 - mable.reg, 41
 - maple.ph, 56
- * **Density deconvolution**
 - mable.decon, 21
- * **Generalized proportional odds model**
 - mable.po, 38
- * **Mixture beta distribution**
 - dmixbeta, 7
 - dtmixbeta, 10
- * **accelerated failure time model**
 - mable.reg, 41
- * **copula density estimation**
 - mable.copula, 17
 - mable.hellcorr, 29
- * **copula**
 - mable.copula, 17
 - mable.hellcorr, 29
- * **datasets**
 - chicken.embryo, 3
 - cosmesis, 7
 - pancreas, 66
 - Vaal.Flow, 71
- * **density estimation**
 - mable.mvar, 33
 - made.mvar, 45
- * **distribution**
 - dmixbeta, 7
 - dtmixbeta, 10
 - mable, 11
 - mable.aft, 15
 - mable.copula, 17
 - mable.group, 27
 - mable.hellcorr, 29
 - mable.mvar, 33
 - mable.ph, 36
 - mable.po, 38
 - mable.reg, 41
 - made.mvar, 45
 - maple.aft, 49
 - maple.ph, 56
 - maple.po, 58
- * **interval censoring**
 - mable.aft, 15
 - mable.ph, 36
 - mable.po, 38
 - mable.reg, 41
 - maple.aft, 49
 - maple.ph, 56
 - maple.po, 58
- * **models**
 - dmixbeta, 7
 - dtmixbeta, 10
 - mable, 11
 - mable.aft, 15
 - mable.group, 27
 - mable.ph, 36
 - mable.po, 38
 - mable.reg, 41
 - maple.aft, 49
 - maple.ph, 56
 - maple.po, 58
- * **multivariate Bernstein polynomial model for copula**
 - mable.copula, 17
 - mable.hellcorr, 29

- * **multivariate Bernstein polynomial model**
 - mable.mvar, 33
 - made.mvar, 45
- * **multivariate**
 - mable.copula, 17
 - mable.hellcorr, 29
 - mable.mvar, 33
 - made.mvar, 45
- * **nonparametric**
 - dmixbeta, 7
 - dtmixbeta, 10
 - mable, 11
 - mable.aft, 15
 - mable.copula, 17
 - mable.group, 27
 - mable.hellcorr, 29
 - mable.mvar, 33
 - mable.ph, 36
 - mable.po, 38
 - mable.reg, 41
 - made.mvar, 45
 - maple.aft, 49
 - maple.ph, 56
 - maple.po, 58
- * **proportional odds model**
 - maple.po, 58
- * **regression**
 - mable.aft, 15
 - mable.ph, 36
 - mable.po, 38
 - mable.reg, 41
 - maple.aft, 49
 - maple.ph, 56
 - maple.po, 58
- * **smooth**
 - dmixbeta, 7
 - dtmixbeta, 10
 - mable, 11
 - mable.aft, 15
 - mable.group, 27
 - mable.ph, 36
 - mable.po, 38
 - mable.reg, 41
 - maple.aft, 49
 - maple.ph, 56
 - maple.po, 58
- * **survival**
 - mable.aft, 15
 - mable.ph, 36
 - mable.po, 38
 - mable.reg, 41
 - maple.aft, 49
 - maple.ph, 56
 - maple.po, 58
- * **univar**
 - mable, 11
 - mable.group, 27
- chicken.embryo, 3
- chpt.exp, 4
- copula2d, 4
- copula2d.cond, 5
- corr.hellinger, 6
- cosmesis, 7
- d2dcop.asym (copula2d), 4
- dcopula (copula2d), 4
- dcopula.cond (copula2d.cond), 5
- dmixbeta, 7
- dmixmbeta, 9
- dtmixbeta, 10
- hellcorr (mable.hellcorr), 29
- mable, 9, 11, 11, 19, 31, 34, 35, 46, 47
- mable.aft, 15, 42, 51
- mable.copula, 17, 31
- mable.ctrl, 12, 15, 18, 20, 21, 24, 26, 27, 31, 32, 34, 36, 39, 42, 44, 46, 50, 52, 54, 56, 59, 69, 72
- mable.decon, 21
- mable.dr, 23
- mable.dr.group, 25
- mable.group, 27, 33
- mable.hellcorr, 29
- mable.ic, 29, 31
- mable.mvar, 19, 31, 33
- mable.ph, 36, 42, 58
- mable.po, 38, 42, 60
- mable.reg, 41
- made.copula, 43
- made.density, 44
- made.mvar, 45
- madem.copula, 47
- madem.density, 48
- maple.aft, 17, 49
- maple.dr, 52

maple.dr.group, 54
maple.ph, 37, 38, 40, 56
maple.po, 40, 58
marginal.p, 61
mvecdf, 62
mvpbeta, 62

optim.gcp, 63
optimable, 13, 14, 34, 35, 46, 47, 64

p2dcop.asym(copula2d), 4
pancreas, 66
pcopula(copula2d), 4
pcopula.cond(copula2d.cond), 5
plot.mable, 67
plot.mable_reg, 68
pmixbeta(dmixbeta), 7
pmixmvbeta(dmixmvbeta), 9
ptmixbeta(dtmixbeta), 10

qcopula.cond(copula2d.cond), 5
qmixbeta(dmixbeta), 7
qtmixbeta(dtmixbeta), 10

r2dcop.asym(copula2d), 4
rcopula(copula2d), 4
rcopula.cond(copula2d.cond), 5
rmixbeta(dmixbeta), 7
rmixmvbeta(dmixmvbeta), 9
rtmixbeta(dtmixbeta), 10

se.coef.dr, 69
summary.mable, 70
summary.mable_reg(summary.mable), 70

umc.mat, 70

Vaal.Flow, 71

weib.gpo, 71