

# Package: pkmon (via r-universe)

September 7, 2024

**Type** Package

**Title** Least-Squares Estimator under k-Monotony Constraint for Discrete Functions

**Version** 1.1

**Date** 2023-08-28

**Author** Jade Giguelay

**Maintainer** Francois Deslandes <francois.deslandes@inra.fr>

**Description** We implement two least-squares estimators under k-monotony constraint using a method based on the Support Reduction Algorithm from Groeneboom et al (2008) <[DOI:10.1111/j.1467-9469.2007.00588.x](https://doi.org/10.1111/j.1467-9469.2007.00588.x)>. The first one is a projection estimator on the set of k-monotone discrete functions. The second one is a projection on the set of k-monotone discrete probabilities. This package provides functions to generate samples from the spline basis from Lefevre and Loisel (2013) <[DOI:10.1239/jap/1378401239](https://doi.org/10.1239/jap/1378401239)>, and from mixtures of splines.

**License** CC BY 4.0

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2023-08-28 09:10:02 UTC

## Contents

pkmon-package . . . . .	2
BaseNorm . . . . .	5
Delta . . . . .	6
estMonotone . . . . .	7
kKnot . . . . .	8
pEmp . . . . .	9
Spline . . . . .	10

<b>Index</b>	<b>12</b>
--------------	-----------

---

pkmon-package	<i>Least-squares estimator under k-monotony constraint for discrete functions</i>
---------------	---

---

## Description

Description: This package implements two least-squares estimators under k-monotony constraint using a method based on the Support Reduction Algorithm from Groeneboom et al (2008). The first one is a projection estimator on the set of k-monotone discrete functions. The second one is a projection on the set of k-monotone discrete probabilities. This package provides functions to generate samples from the spline basis from Lefevre and Loisel (2013), and from mixtures of splines.

## Author(s)

Jade Giguelay

## References

- Giguelay J. (2016) Estimation of a discrete distribution under k-monotony constraint, *in revision*, (arXiv:1608.06541)
- Lefevre C., Loisel S. (2013) <DOI:10.1239/jap/1378401239> On multiply monotone distributions, continuous or discrete, with applications, *Journal of Applied Probability*, **50**, 827–847.
- Groeneboom P., Jongbloed G. Wellner J. A. (2008) <DOI:10.1111/j.1467-9469.2007.00588.x> The Support Reduction Algorithm for Computing Non-Parametric Function Estimates in Mixture Models, *Scandinavian Journal of Statistics*, **35**, 385–399

## See Also

[pMonotone](#), [fMonotone](#)

## Examples

```
#####
# Example 1
# one triangular function  $T_j=Q_j^2$ , for  $j=\text{supp}=20$  and for  $k=2$  and  $k=3$ 
n=30;
k1=2;
k2=3;
l=2;
supp=20;
p=dSpline(supp, k=1);
X=rSpline(n=n, supp, k=1);
ptilde=pEmp(X);
phat1=pMonotone(ptilde$freq, k=k1);
phat2=pMonotone(ptilde$freq, k=k2);

x.limits=c(0, max(supp+1, phat1$Spi+1, phat2$Spi+1));
```

```

y.limits=range(p, ptilde$freq, phat1$p, phat2$p);

plot(NULL, xlim=x.limits, ylim=y.limits, xlab="Counts", ylab="Frequencies");
points(0:supp, p, pch=16, col=1, lwd=2);
points(ptilde$supp, ptilde$freq, pch=4, col=2, lwd=2);
points(0:max(phat1$Spi), phat1$p, pch=8, col=3, lwd=2);
points(0:max(phat2$Spi), phat2$p, pch=2, col=4, lwd=2);
legend("topright", pch=c(16, 4, 8, 2), col=c(1, 2, 3, 4),
      legend=c("p", expression(tilde("p")), expression(hat("p")*" - k = 2"),
      expression(hat("p")*" - k = 3")));

#####
# Example 2
# mixture of 3 splines  $Q_j^3$  and for  $k=4$  and  $k=3$ 
n=30;
k1=4;
k2=3;
l=3;
supp=c(5, 10, 20);
prob=c(0.5, 0.3, 0.2);
p=dmixSpline(supp, k=l, prob=prob);
X=rmixSpline(n=n, supp, k=l, prob=prob);
ptilde=pEmp(X);
phat1=pMonotone(ptilde$freq, k=k1);
phat2=pMonotone(ptilde$freq, k=k2);

x.limits=c(0, max(supp+1, phat1$Spi+1, phat2$Spi+1));
y.limits=range(p, ptilde$freq, phat1$p, phat2$p);

plot(NULL, xlim=x.limits, ylim=y.limits, xlab="Counts", ylab="Frequencies");
points(0:max(supp), p, pch=16, col=1, lwd=2);
points(ptilde$supp, ptilde$freq, pch=4, col=2, lwd=2);
points(0:max(phat1$Spi), phat1$p, pch=8, col=3, lwd=2);
points(0:max(phat2$Spi), phat2$p, pch=2, col=4, lwd=2);
legend("topright", pch=c(16, 4, 8, 2), col=c(1, 2, 3, 4),
      legend=c("p", expression(tilde("p")), expression(hat(p)* " - k = 4"),
      expression(hat(p)* " - k = 3")));

#####
# Example 3
# Poisson density
n=30;
k1=2;
k2=3;
supp=10;
p=dpois(0:supp, lambda=1);
X=rpois(n, lambda=1);
ptilde=pEmp(X);
phat1=pMonotone(ptilde$freq, k=k1);
phat2=pMonotone(ptilde$freq, k=k2);

x.limits=c(0, max(supp, phat1$Spi+1, phat2$Spi+1));
y.limits=range(p, ptilde$freq, phat1$p, phat2$p);

```

```

plot(NULL, xlim=x.limits, ylim=y.limits, xlab="Counts", ylab="Frequencies");
points(0:max(supp), p, pch=16, col=1, lwd=2);
points(ptilde$supp, ptilde$freq, pch=4, col=2, lwd=2);
points(0:max(phat1$Spi), phat1$p, pch=8, col=3, lwd=2);
points(0:max(phat2$Spi), phat2$p, pch=2, col=4, lwd=2);
legend("topright", pch=c(16, 4, 8, 2), col=c(1, 2, 3, 4),
      legend=c("p", expression(tilde("p")), expression(hat(p)* " - k = 2"),
              expression(hat(p)* " - k = 3")));

## Not run:
#####
# Simulation for comparing ptilde and pHat (p is 3-monotone, k=3)
#
#OUTPUT
#
# cvge : percentage of non-convergence of the algorithm
# r.emp : L2-risk for the empirical estimator
# r.Hat : L2-risk for the estimator under k-monotony constraint

nSim=500;

n=30;
k=3;
l=3;
supp=20;
p=dSpline(supp, k=1);

result <- matrix(nrow=nSim,ncol=3);
dimnames(result)[[2]] <- c("cvge", "r.emp", "r.Hat");
for (i in 1:nSim) {
  X=rSpline(n=n, supp, k=1);
  ptilde=pEmp(X);
  phat=pMonotone(ptilde$freq, k=k);
  m <- max(supp+1,length(ptilde$freq)+1,phat$Spi+1)
  pV=c(p,rep(0,m-length(p)))
  pHat=c(phat$p,rep(0,m-length(phat$p)))
  ptilde=c(ptilde$freq,rep(0,m-length(ptilde$freq)))
  result[i,] <- c(phat$cvge,sum((pV-ptilde)**2),
                sum((pV-pHat)**2))
}

apply(result,2,mean)

#Example with set.seed(0)
#cvge      r.emp      r.Hat
#0.00000000 0.030682552 0.004984899

## End(Not run)

```

---

BaseNorm	<i>Normalized spline basis</i>
----------	--------------------------------

---

**Description**

Computes the k-monotone discrete splines from Lefevre and Loisel (2013).

**Usage**

```
BaseNorm(k, J)
```

**Arguments**

k	Degree of monotony
J	maximum support of the splines

**Value**

matrix  $Q$  with  $J+1$  rows and  $J+1$  columns with  $Q(i, j) = Q_j^k(i - 1) = C_{j-i+k-1}^{k-1}$ , where  $C$  represents the binomial coefficient.

**Author(s)**

Jade Giguelay

**References**

Giguelay, J., (2016), Estimation of a discrete distribution under k-monotony constraint, *in revision*, (arXiv:1608.06541)

Lefevre C., Loisel S. (2013) <DOI:10.1239/jap/1378401239> On multiply monotone distributions, continuous or discrete, with applications, *Journal of Applied Probability*, **50**, 827–847.

**See Also**

[rSpline](#), [dSpline](#), [rmixSpline](#), [dmixSpline](#)

**Examples**

```
# Computing 3-monotone splines with maximum support 8
Q=BaseNorm(3, 8)
matplot(Q, type="l", main="3-monotone splines with maximum support 8");
```

---

Delta

*Discrete laplacian*

---

### Description

Computes the laplacians of a discrete function

### Usage

Delta(k, L, p)

### Arguments

k	Maximum order of the laplacian
L	Support of the function
p	Discrete function represented as a vector

### Value

Returns a matrix with the laplacians  $(-1)^j \Delta^j(p(l))$  of vector  $p$  for  $j$  in  $1, \dots, k$  and  $l$  in  $0, \dots, L$ .

### Author(s)

Jade Giguelay

### References

Knopp K. (1925), <DOI:10.1007/BF01479598> Mehrfach monotone Zahlenfolgen, *Mathematische Zeitschrift*, **22**, 75–85

Giguelay, J., (2016), Estimation of a discrete distribution under k-monotony constraint, *in revision*, (arXiv:1608.06541)

### See Also

[kKnot](#)

### Examples

```
p=dSpline(k=3, supp=20)
M=Delta(3, 20, p)
```

estMonotone

*Estimators of discrete probabilities under k-monotony constraint***Description**

Estimators of discrete probabilities under k-monotony constraint. Estimation can be done on the set of k-monotone functions or on the set of k-monotone probabilities.

**Usage**

```
pMonotone(ptild, t.zero = 1e-10, t.P = 1e-08, max.sn = NULL, k, verbose = FALSE)
fMonotone(ptild, t.zero = 1e-10, t.P = 1e-08, max.sn = NULL, k, verbose = FALSE)
```

**Arguments**

ptild	Empirical estimator
t.zero	Threshold for the precision of the directionnal derivatives. (see OUTPUT below)
t.P	Threshold for the precision on the stopping criterion. (see OUTPUT below)
max.sn	The maximum support for the evaluation of the estimator
k	Degree of monotony
verbose	if TRUE, print for each iteration on the maximum support : pi, Psi and sumP (see OUTPUT below)

**Details**

The thresholds t.P and t.zero are used for the precision in the algorithm : in Step one (See REFERENCES below) the algorithm computes the directionnal derivatives of the current estimator and stops if all the directionnal derivarives are null that is to say if they are smaller than t.zero. In Step two (See REFERENCES below) the algorithm computes a stopping criterion and stops if and only if the stopping criterion is verified that is to say if some quantities are non-negative that is to say bigger than -t.P.

**Value**

cvge	cvge = 0 if the criterion Psi decreases with the support of pi. cvge = 1 if Psi increases. cvge = 2 if maximum number of iterations reached
Spi	Support of the positive measure pi at the last iteration
pi	Values of the positive measure pi at the last iteration
p	Values of pHat
Psi	Scalar value of the criterion to be minimised
sumP	sum(pHat) at convergence
history	Data frame with components
L	The maximum of the support of pi
Psi	Value of the criterion for the value L
SumP	Value of sum(pHat)

**Author(s)**

Jade Giguelay

**References**

Giguelay, J., (2016), Estimation of a discrete distribution under k-monotony constraint, *in revision*, (arXiv:1608.06541)

Groeneboom P., Jongbloed G. Wellner J. A. (2008) <DOI:10.1111/j.1467-9469.2007.00588.x> The Support Reduction Algorithm for Computing Non-Parametric Function Estimates in Mixture Models, *Scandinavian Journal of Statistics*, **35**, 385–399

**See Also**

[pEmp](#), [BaseNorm](#)

**Examples**

```
x=rSpline(n=50, 20, k=4)
ptild=pEmp(x);
res=pMonotone(ptild$freq, k=4)
```

---

kKnot

*k-Knot*


---

**Description**

k-Knots of a discrete function.

**Usage**

```
kKnot(p, k)
```

**Arguments**

p	Vector
k	Degree of the knots

**Details**

An integer  $i$  is a  $k$ -knot of  $p$  if  $\Delta^k p(i) > 0$ , where  $\Delta^k$  is the  $k$ -th Laplacian of the sequence  $p$ .

**Value**

Vector with the  $k$ -knots of  $p$ .

**Author(s)**

Jade Giguelay



**References**

Knopp K. (1925), <DOI:10.1007/BF01479598> Mehrfach monotone Zahlenfolgen, *Mathematische Zeitschrift*, **22**, 75–85

Giguelay, J., (2016), Estimation of a discrete distribution under k-monotony constraint, *in revision*, (arXiv:1608.06541)

**See Also**

[Delta](#)

**Examples**

```
p=dmixSpline(c(5, 10, 20), k=3, c(0.5, 0.25, 0.25))
knots=kKnot(p, 3)
```

---

pEmp

*Empirical estimator of a discrete function*

---

**Description**

Empirical estimator of a discrete function

**Usage**

pEmp(X)

**Arguments**

X                      A random sample from a discrete probability.

**Details**

The empirical estimator is defined as  $p(j) = \frac{1}{n} \sum_{i=1}^n \mathbf{1}_{x_j=j}$ .

**Value**

support	The points of the support of the estimator
count	The counts of the sample
freq	The normalized counts

**Author(s)**

Jade Giguelay

**Examples**

```
x=rpois(100, lambda=0.3)
ptild=pEmp(x)
```

---

Spline

*Random generation and distribution function of k-monotone densities*

---

### Description

Random generation and distribution function for the spline of the basis from Lefevre and Loisel (2013), and mixtures of splines.

### Usage

```
rSpline(n=1, supp, k)
dSpline(supp, k)
rmixSpline(n=1, supp, k,prob)
dmixSpline(supp, k, prob)
```

### Arguments

supp	Support of the spline, or vector of the supports of the splines for the mixture of splines
n	Number of random values to return
k	Degree of monotony
prob	Vector of probabilities for the mixture of splines

### Details

See [BaseNorm](#) for details on the spline basis.

### Value

rSpline and rmixSpline generates random deviates from the splines and mixtures of splines.  
dSpline and dmixSpline gives the distribution function.

### Author(s)

Jade Giguelay

### References

Giguelay, J., (2016), Estimation of a discrete distribution under k-monotony constraint, *in revision*, (arXiv:1608.06541)  
Lefevre C., Loisel S. (2013) <DOI:10.1239/jap/1378401239> On multiply monotone distributions, continuous or discrete, with applications, *Journal of Applied Probability*, **50**, 827–847.

### See Also

[pEmp](#)

**Examples**

```
x=rSpline(n=100, 20, 3)
p=dSpline(20, 3)
xmix=rmixSpline(n=100, c(5, 20), 3, c(0.5, 0.5))
pmix=dmixSpline(c(5, 20), 3, c(0.5, 0.5))
par(mfrow=c(1, 2))
hist(x, freq=FALSE)
lines(p, col="red")
hist(xmix, freq=FALSE)
lines(pmix, col="red")
```

# Index

## \* **package**

pkmon-package, 2

BaseNorm, 5, 8, 10

Delta, 6, 9

dmixSpline, 5

dmixSpline (Spline), 10

dSpline, 5

dSpline (Spline), 10

estMonotone, 7

fMonotone, 2

fMonotone (estMonotone), 7

kKnot, 6, 8

pEmp, 8, 9, 10

pkmon (pkmon-package), 2

pkmon-package, 2

pMonotone, 2

pMonotone (estMonotone), 7

rmixSpline, 5

rmixSpline (Spline), 10

rSpline, 5

rSpline (Spline), 10

Spline, 10