

Package: scpropreg (via r-universe)

May 15, 2026

Type Package

Title Simplicially Constrained Regression Models for Proportions

Version 1.2

Date 2026-04-15

Author Michail Tsagris [aut, cre]

Maintainer Michail Tsagris <mtsagris@uoc.gr>

Depends R (>= 4.0)

Imports nnsolve, Rfast, Rfast2

Description Simplicially constrained regression models for proportions in both sides. The constraint is always that the betas are non-negative and sum to 1. References: Iverson S.J., Field C., Bowen W.D. and Blanchard W. (2004) ``Quantitative Fatty Acid Signature Analysis: A New Method of Estimating Predator Diets". Ecological Monographs, 74(2): 211-235. <doi:10.1890/02-4105>.

License GPL (>= 2)

NeedsCompilation no

Repository <https://cran.r-universe.dev>

Date/Publication 2026-04-15 11:30:48 UTC

RemoteUrl <https://github.com/cran/scpropreg>

RemoteRef HEAD

RemoteSha 89c661226127c1bd2e9484cc224790a6c8340b8e

Contents

scpropreg-package	2
Positive and unit sum constrained least squares	2
QFASA diet estimates for many predators using various distances	3
QFASA diet estimates using various distances	5
Index	7

scpropreg-package

Simplicially Constrained Regression Models for Proportions

Description

Simplicially Constrained Regression Models for Proportions. The constraint is always that the betas are non-negative and sum to 1.

Details

Package: scpropreg
Type: Package
Version: 1.2
Date: 2026-04-15

Maintainers

Michail Tsagris <mtsagris@uoc.gr>.

Author(s)

Michail Tsagris <mtsagris@uoc.gr>

References

Iverson Sara J., Field Chris, Bowen W. Don and Blanchard Wade (2004) Quantitative Fatty Acid Signature Analysis: A New Method of Estimating Predator Diets. *Ecological Monographs*, 74(2): 211-235.

Positive and unit sum constrained least squares

Positive and unit sum constrained least squares

Description

Positive and unit sum constrained least squares.

Usage

pcls(y, x)
mpcls(Y, x)

Arguments

y	The response variable. For the pcls() a numerical vector with observations, but for the mpcls() a numerical matrix, where each columns sums to 1.
Y	The response variable, a numerical matrix, where each column sums to 1.
x	A matrix with independent variables, the design matrix.

Details

The constraint is that all beta coefficients are positive and sum to 1. That is $\min \sum_{i=1}^n (y_i - \mathbf{x}_i \top \boldsymbol{\beta})^2$ such that $0 \leq \beta_j \leq 1$ and $\sum_{j=1}^d \beta_j = 1$. The pcls() function performs a single regression model, whereas the mpcls() function performs a regression for each column of y. Each regression is independent of the others.

Value

A list including:

coefficients	A numerical matrix with the positively constrained beta coefficients.
value	A numerical vector with the mean squared error.

Author(s)

Michail Tsagris.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr>.

See Also

[kld](#), [mkld](#)

Examples

```
x <- matrix(runif(30 * 8), ncol = 30)
x <- t( x / rowSums(x) )
y <- runif(30)
y <- y / sum(y)
pcls(y, x)
```

QFASA diet estimates for many predators using various distances

QFASA diet estimates for many predators using various distances

Description

QFASA diet estimates for many predators using various distances.

Usage

```

mkld(Y, x, tol = 1e-8, maxit = 50000, alpha = 0.1)
mait(Y, x, tol = 1e-8, maxit = 50000, alpha = 0.01)
mlsq(Y, x, tol = 1e-8, maxit = 50000, alpha = 0.01)
mlr(Y, x, tol = 1e-8, maxit = 100)

```

Arguments

Y	The response variable, a matrix with values between 0 and 1 that sum to 1. For some functions, zero values are allowed. Every column corresponds to the food composition of a predator. The column-wise sums are equal to 1.
x	A matrix with independent variables, values between 0 and 1. Each column contains a prey's diet. The column-wise sums are equal to 1.
tol	The tolerance value to terminate the algorithm.
maxit	The maximum iterations allowed.
alpha	The step-size parameter of the fixed points iteration algorithm. This is similar to the η parameter in the gradient descent algorithm.

Details

The function estimates the betas that minimize a distance. The fitted values are linear constraints of the observed xs. The constraint is that all beta coefficients are positive and sum to 1. That is $\hat{y}_i = \sum_{j=1}^d x_{ij}\beta_j$ such that $0 \leq \beta_j \leq 1$ and $\sum_{j=1}^d \beta_j = 1$.

Value

A list including:

coefficients	A numerical matrix with the positively constrained beta coefficients.
value	A numerical vector with the value of the objective function.
iters	The number of iterations required until termination of the algorithm.

Author(s)

Michail Tsagris.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr>.

References

Iverson Sara J., Field Chris, Bowen W. Don and Blanchard Wade (2004) Quantitative Fatty Acid Signature Analysis: A New Method of Estimating Predator Diets. *Ecological Monographs*, 74(2): 211-235.

See Also

[kld](#), [mpcls](#)

Examples

```
x <- matrix(runif(30 * 6), ncol = 30)
x <- t( x / rowSums(x) )
Y <- matrix(runif(30 * 10), ncol = 30)
Y <- t( Y / rowSums(Y) )
mkld(Y, x)
```

QFASA diet estimates using various distances

QFASA diet estimates using various distances

Description

QFASA diet estimates using various distances.

Usage

```
klld(y, x, tol = 1e-8, maxit = 50000, alpha = 0.1)
ait(y, x, tol = 1e-8, maxit = 50000, alpha = 0.01)
lsq(y, x, tol = 1e-8, maxit = 50000, alpha = 0.01)
jsd(y, x, tol = 1e-8, maxit = 300000, alpha = 0.01)
lr(y, x, tol = 1e-8, maxit = 100)
```

Arguments

y	The response variable. The predator's food composition. A vector with values between 0 and 1 that sum to 1. For some functions, zero values are allowed.
x	A matrix with independent variables, values between 0 and 1. Each column contains a prey's diet. The column-wise sums are equal to 1.
tol	The tolerance value to terminate the algorithm.
maxit	The maximum iterations allowed.
alpha	The step-size parameter of the fixed points iteration algorithm. This is similar to the η parameter in the gradient descent algorithm.

Details

The function estimates the betas that minimize a distance. The fitted values are linear constraints of the observed xs. The constraint is that all beta coefficients are positive and sum to 1. That is $\hat{y}_i = \sum_{j=1}^d x_{ij} \beta_j$ such that $0 \leq \beta_j \leq 1$ and $\sum_{j=1}^d \beta_j = 1$.

Value

A list including:

coefficients	A numerical matrix with the positively constrained beta coefficients.
value	A numerical vector with the value of the objective function.
iters	The number of iterations required until termination of the algorithm.

Author(s)

Michail Tsagris.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr>.

References

Iverson Sara J., Field Chris, Bowen W. Don and Blanchard Wade (2004) Quantitative Fatty Acid Signature Analysis: A New Method of Estimating Predator Diets. *Ecological Monographs*, 74(2): 211-235.

See Also

[mkld](#), [pcls](#)

Examples

```
x <- matrix(runif(30 * 8), ncol = 30)
x <- t( x / rowSums(x) )
y <- runif(30)
y <- y / sum(y)
kld(y, x)
ait(y, x)
lsq(y, x)
lr(y, x)
```

Index

- ait (QFASA diet estimates using various distances), [5](#)
- jsd (QFASA diet estimates using various distances), [5](#)
- kld, [3](#), [4](#)
- kld (QFASA diet estimates using various distances), [5](#)
- lr (QFASA diet estimates using various distances), [5](#)
- lsq (QFASA diet estimates using various distances), [5](#)
- mait (QFASA diet estimates for many predators using various distances), [3](#)
- mkld, [3](#), [6](#)
- mkld (QFASA diet estimates for many predators using various distances), [3](#)
- mlr (QFASA diet estimates for many predators using various distances), [3](#)
- mlsq (QFASA diet estimates for many predators using various distances), [3](#)
- mpcls, [4](#)
- mpcls (Positive and unit sum constrained least squares), [2](#)
- pcls, [6](#)
- pcls (Positive and unit sum constrained least squares), [2](#)
- Positive and unit sum constrained least squares, [2](#)
- QFASA diet estimates for many predators using various distances, [3](#)
- QFASA diet estimates using various distances, [5](#)
- scpropreg-package, [2](#)